



Herbivory and seedling establishment in Pyrenean forests: Influence of micro- and meso-habitat factors on browsing pressure



Aitor Ameztegui^{a,b,*}, Lluís Coll^{a,c}

^a Forest Sciences Centre of Catalonia (CEMFOR-CTFC), Ctra. Sant Llorenç de Morunys km. 2, E-25280 Solsona, Catalonia, Spain

^b Centre for Forest Research (CEF), Département des Sciences Biologiques, Université du Québec à Montréal, Montréal, Québec H3C 3P8, Canada

^c CREA, Cerdanyola del Vallès 08193, Spain

ARTICLE INFO

Article history:

Received 26 November 2014

Received in revised form 19 January 2015

Accepted 22 January 2015

Keywords:

Browsing

Mortality

Ungulates

Livestock

Seedlings

Pyrenees

ABSTRACT

Browsing damage is among the most determinant factors that limit the establishment of tree seedlings in forests. In some areas, this process leads to massive mortalities that can reduce or even completely prevent the regeneration of some tree species. Mediterranean mountain forests have undergone during the last decades important changes in land-uses that have significantly altered the type and abundance of herbivore populations. In this study we assessed the impact of current grazing conditions in forest regeneration using a set of experimental plantations established in the Eastern Pyrenees in areas visited by domestic livestock (cattle and horses) and wild ungulates (mainly roe deer and chamois). We analyzed during 4 years the role of seedling species and size, mesohabitat (elevation and type of forest cover) and microhabitat (herbaceous cover, distance to shrub, and light availability) on the browsing-induced mortality of more than 500 seedlings of *Pinus sylvestris*, *Pinus uncinata*, *Betula pendula* and *Abies alba*, the most common tree species in the study area. Browsing-induced mortality for the three conifer species was much lower (<15%) than the one observed for *B. pendula* (>40%) and depended on both microhabitat – mainly on the distance to protective elements such as shrubs; and mesohabitat, with an interaction between the elevational belt (site) and the type of forest cover (gaps vs. understory). In the subalpine belt, browsing on *A. alba* and *P. uncinata* was higher during summer at plots located in the forest understory whereas, during winter, it was higher at plots located in gaps. The study shows that both mesohabitat and microhabitat can exert an effect on the patterns of plant damage by herbivores, providing useful information to adapt forest management in areas particularly exposed to them.

© 2015 Elsevier B.V. All rights reserved.

1. Introduction

Forests regeneration is limited by several environmental constraints. Among them, browsing by large herbivores is often one of the most determinant factors, particularly in mountain areas, where tree establishment and growth are already constrained by unfavorable environmental conditions (Ameztegui and Coll, 2013). In the last decades, many European mountains have undergone major changes in the type, abundance and behavior of their main herbivore populations – including both wild and domestic animals – as a consequence of the profound socioeconomic changes that have affected these systems (Casals et al., 2009). For instance, the Pyrenees have recently changed from a transhumant system – in which sheep for wool production were favored against

cattle, and there was an extensive utilization of most available food sources – to a system in which the overall livestock pressure is notably lower, and beef cattle and breeding mares have partially substituted sheep herds (García-Ruiz and Lasanta, 1990; Lasanta, 1990). Under this new organization of the landscape, intensive grazing pressures concentrate in very limited areas (mainly sub-alpine pastures), whereas many pastures that were suitable for sheep are now unused (Balcells, 1983). The abandonment of these lands together with a reduction of logging pressure on forests have led to important forest expansion processes and to a generalized increase in canopy closure (Ameztegui et al., 2010; Poyatos et al., 2003). Wild ungulates such as roe deer (*Capreolus capreolus*) or chamois (*Rupicapra pyrenaica*) find in these dense forests a suitable habitat and have sharply increased their populations in the last decades (Apollonio et al., 2010; Putman et al., 2011).

The progressive increase in ungulate populations during the last decades has already caused some concern due to the consequences it could entail for the regeneration of European temperate forests (see, for example Motta, 1996; Putman, 1996 or Bugmann and

* Corresponding author at: Centre for Forest Research (CEF), Département des Sciences Biologiques, Université du Québec à Montréal, C.P. 8888, Succ. Centre-Ville, Montréal, Québec H3C 3P8, Canada. Tel.: +1 (514) 987 3000/4866.

E-mail address: ameztegui@gmail.com (A. Ameztegui).

Weisberg, 2003). At high stocking densities, herbivory can even completely prevent regeneration of some species and lead to a reduction of plant species diversity (Ammer, 1996; Danell, 2003; Reimoser, 1996). Changes in herbivore populations – both in livestock and wildlife – can severely affect the patterns of seedling consumption and lead to shifts in forest composition on the mid-term (Gill, 1992; Motta, 1996; Senn and Suter, 2003), and patterns of seedling predation by large herbivores can themselves be affected by other environmental factors (Weisberg and Bugmann, 2003). Available tree species, forest structure, type of habitat, period of grazing or the amount of alternative food sources, among others, affect how frequently and intensely herbivores browse young trees (Liss, 1988; Mayer, 2005a; Mayer et al., 2003; Reimoser, 1996). Yet, we are still far from a complete understanding of the combined effect that these environmental factors can exert on browsing patterns (Bergström and Edenius, 2003; Kräuchi et al., 2000).

In this study, we aim at shedding some light on the role that the complex ungulate-habitat interactions play on the dynamics of mountain forests. For this purpose, we planted seedlings of the main tree species in the Eastern Pyrenees in an area grazed both by livestock and wild ungulates, and under contrasting environmental conditions. By planting even-aged seedlings in areas with non-modified browsing conditions, we wanted to assess the effect of specific seedling selection by herbivores (Kupferschmid et al., 2014). Mortality of seedlings due to browsing, although often non-visible, can be more determinant on forest dynamics than mortality of saplings, but the later has been more studied because the damages are more evident and easier to determine (Reimoser, 2003; Senn and Suter, 2003). More specifically, the study aimed to (i) assess the impact of current grazing conditions (both of livestock and wild ungulates) in forest establishment; (ii) detect if some tree species were preferred by herbivores over others; (iii) determine how variations in meso-habitat (elevation and type of forest cover) and microhabitat (herbaceous cover, distance to shrub, light availability) may engender different patterns of plant consumption by herbivores; and (iv) discuss the potential implications of the observed patterns for forest dynamics. We expect a highly selective predation on silver fir and birch, which can prevent the diversification of pine forest described elsewhere in the absence of management (Navarro-González et al., 2013). At the meso-habitat scale, we expect livestock to cause less damage than wild ungulates at forest gaps; whereas at the micro-habitat scale, the predation rate is likely to be positively affected by herb abundance and negatively by shrubs.

2. Material and methods

2.1. Study area and species

We conducted the experiment at two sites with different elevations in the northern slopes of the Serra del Cadí, a Pyrenean mountain range in the Cadí-Moixeró Natural Park (42°17'N; 1°42'E). The Sant Romà site corresponds to a montane pine forest located at 1500 m a.s.l. (near the *Pinus sylvestris* – *Pinus uncinata* transition zone), while the Prat d'Aguiló site is a typical subalpine forest located close to the optimum elevational distribution for *P. uncinata* (around 2000 m a.s.l.; Fig. 1). The forest in Sant Romà was dominated by *P. sylvestris*, with some *P. uncinata* and *Betula pendula* individuals, and the main species in the understory were common box (*Buxus sempervirens* L.) and common juniper (*Juniperus communis* L.). Prat d'Aguiló forests were dominated by *P. uncinata* and the shrub layer was almost exclusively composed by *J. communis*.

Despite being a protected area, herds of cows and horses graze every summer in both sites (Table 1), since the authorities have

promoted animal husbandry as a major source of income for local communities. The area also hosts important populations of chamois (*R. pyrenaica*), and roe deer (*C. capreolus*), with a lower presence of wild boar (*Sus scrofa*), and red deer (*Cervus elaphus*).

We used for this study the four most common tree species in the study area: Scots pine (*P. sylvestris* L.), mountain pine (*P. uncinata* Ram. ex DC), silver fir (*Abies alba* Mill.) and silver birch (*B. pendula* Roth). These species have different ecological requirements in terms of tolerance to shade, drought and cold temperatures (see Ameztegui and Coll, 2013 for further details) but they can coexist between 1600 and 2000 m a.s.l., where they constitute the montane–subalpine ecotone.

2.2. Experimental design

A total of 1152 seedlings of the study species were planted in the early summer of 2008. Seedlings were distributed along 24 plots (12 at the montane site and 12 at the subalpine), located at less than 300 m from grasslands highly frequented by livestock (Fig. 1). At each site, half of the plots (6) were established in the forest understory and the other half (6) in naturally occurring gaps, whose size ranged between 150 and 350 m². Each plot contained 48 seedlings (12 per species) randomly distributed in a grid pattern, and seedlings were planted at least 1 m one from another to avoid above- and below-ground interaction. All the seedlings were two years old at the moment of plantation, and had been grown in a local nursery (Forestal Catalana, Pobla de Lillet, Spain) from seeds collected in neighboring forests, i.e., seed source, nursery, and plantation area were all inside the same provenance region (Alfía et al., 2005). To ensure enough seedling survival to assess the role of climate and microsite on seedling performance (see Ameztegui and Coll, 2013), half of the seedlings in each plot were protected from browsing with an individual protector (90 cm height and 33 cm of diameter) with a mesh net of 20 × 20 mm (Nortène, Lille, France). The remaining half ($n = 576$) was left unprotected and constitute the sample of this study, since they allowed us to assess for the effects of browsing by large herbivores on seedling performance.

2.3. Presence of herbivores in the study area

Both sites are highly frequented by livestock during summer, when a total of 95 cows and 60 mares – i.e. 124 adult livestock units – take advantage of the pastures located close to the experimental sites under a free-range, extensive system. During summer, the high presence of domestic cattle forces most wildlife to take refuge in the rocky areas located above the main grazing areas. On the contrary, wild ungulates are the unique large herbivores in the study area during winter, when livestock is stabled in the valley. Although there are no recent censuses of wild ungulates in the study area, the populations of roe deer, chamois, red deer and wild boar have increased in the last decades (DGMNB, 2014). The study area is located inside the Cadí National Game Reserve, created in 1966 to protect a population of 78 specimens of chamois. The last census of this species performed in 1999 revealed that the population had risen up to 2551 individuals (5.4 chamois km⁻²).

2.4. Seedling monitoring and characterization of the habitat

Individual seedling damage and mortality were monitored at the beginning and at the end of each growing season during four years since the moment of plantation. The likely cause of damage or death was recorded when apparent as whether due to (i) herbivores (browsing, trampling or other type of physical damage caused by browsers); (ii) lack of adaptation (i.e., showing




Fig. 1. Location of the study area inside the (a) Iberian Peninsula; (b) the Pyrenees and (c) the Cadí-Moixerò Natural Park. Stars in (b) and (c) indicate the approximate location of the study plots.

symptoms of drought or cold sensitivity such as partially or totally brown leaves); and (iii) random causes (tree or stone falls, downhill creep). We did not find plants severely affected by other biotic agents such as insects or fungi. We defined browsing-induced mortality as the proportion of the seedlings that died within the study period, and for which the apparent cause of death were herbivores. We also monitored the recovery of those browsed seedlings that did not die immediately, and we computed lethality as the proportion of seedlings that, having been browsed, ended up dying. We described the environment in which seedlings were growing at two spatial scales: mesohabitat (at the plot level) and microhabitat (at the individual level). Mesohabitat was described through the elevational stages of vegetation (montane vs. sub-alpine) and the type of forest cover (understory vs. gap). On the other hand, microhabitat was characterized by light availability, percentage of herbaceous cover and distance to the nearest shrub, measured for every planted seedling (Table 1). Light availability was determined, for each seedling, as percentage of the above canopy photosynthetic photon flux density (%PPFD). Measurements were taken using two Li-190SA quantum sensors (LI-COR, Lincoln, NE, USA) in paired mode, i.e. one of the sensors was placed

at the top of each seedling and the other in an adjacent open area (following Parent and Messier, 1996). The percentage of herbaceous cover surrounding each plant was visually estimated – to the nearest 5% – by placing an 80 × 80 cm reticulated grid centered on each seedling. Distance from each seedling to the nearest shrub – defined as a woody plant with multiple stems arising from the base – was measured with a tape meter.

2.5. Data analyses

For each species, survival function curves were constructed for the unprotected plants ($n = 576$), using Kaplan–Meier estimates based on data of browsed-induced mortality along the study period. This kind of analyses allow for assessment of the occurrence of an event (in this case, the mortality due to browsing) while taking into account censored data, i.e. observations for which the information about their survival time is incomplete, whether because they are alive at the end of the observation period or because they quit the study before completion (in this case, seedlings whose mortality was not related to browsing). Mantel–Cox log-rank tests were used to determine significant differences in the survival

Table 1
Main abiotic and biotic characteristics of the studied stands at the meso-habitat and micro-habitat scale.

	Montane site	Subalpine site
<i>Meso-habitat variables</i>		
Latitude (N)/longitude (E)	42°19'/1°43'	42°18'/1°42'
Elevation (m a.s.l.)	1550	1955
Aspect/slope (degrees)	NE/39	NE/53
Bedrock	Limestone	Limestone
Mean rooting depth (cm)	57.0 ± 9.3	52.5 ± 15.5
Mean annual/summer temperature (°C)	7.4/14.8	4.9/11.7
Total annual/summer precipitation (mm)	992/271	1118/327
Length of the growing season ^a (days)	194	147
Mean summer maximum temperature (°C)	21.0	17.3
Mean winter minimum temperature (°C)	−3.4	−4.6
Dominant species ^b	Ps, Pu, Bp	Pu, Aa, Bp
Main browsing period	June–September	
Livestock units ^c	124 (95 beef cattle + 60 horses)	
<i>Micro-habitat variables</i>		
Light availability (%PPFD)	19.3 ± 9.8 [2.8–55.1]	23.9 ± 13.3 [5.4–58.6]
Herbaceous cover (%)	29.2 ± 19.8 [0–85]	43.0 ± 29.1 [0–100]
Distance to nearest shrub (cm)	59.8 ± 52.5 [10–400]	117.5 ± 82.2 [20–450]

Values for light availability, herbaceous cover, and distance to nearest shrub are mean ± SD. Values in brackets are minimum and maximum observed values.

^a Calculated as in Körner and Paulsen (2004).

^b Listed for each site in decreasing order of dominance: Pu: *Pinus uncinata*; Aa: *Abies alba*; Ps: *Pinus sylvestris*; Bp: *Betula pendula*.

^c Calculated as defined by the European Commission (http://epp.eurostat.ec.europa.eu/statistics_explained/index.php/Glossary:LSU).

curves between sites (montane vs. subalpine) and types of forest cover (gap vs. understory).

To assess the effect of meso- and micro-habitat factors on seedling mortality we computed browsing-induced probability of mortality for every species and plot, using the value of the Kaplan–Meier estimators at the end of the study period. We assessed the effect of mesohabitat factors (elevation and forest cover) on browsing mortality via a split-plot ANOVA in which elevation was considered as the main factor and type of forest cover as the subplot factor. Exploratory analyses showed that the effect of browsing on seedling mortality varied with species, elevation and forest cover, and we found a significant three-way interaction. Hence, to facilitate interpretation of the results, they are presented using within-species analyses. We split the data into summer and winter seasons to test for potential effects of the main type of browser (wild ungulates vs. livestock) on the patterns of mortality. One-sample *t*-tests with Bonferroni corrections were performed to test significant deviations from zero of the mortality values.

To test the effect of micro-habitat on the browsing-induced mortality, each microsite explanatory variable (light availability, herbaceous cover and distance to shrubs) was analyzed separately for each species using logistic models:

$$y = \frac{e^z}{1 + e^z}$$

where *y* is the probability of browsing-induced mortality for an individual sapling, and *z* a parameter to estimate. To allow for non-linear relationships between the response and independent variable, we tested linear, power and exponential formulations for *z*. The procedure used to test the role of an independent variable on seedling performance was the same for all the models, and was based on a likelihood approach. The best of the 3 formulations (linear, power and exponential) was compared to a null model using Δ AICc, the difference in corrected Akaike Information Criterion (Burnham and Anderson, 2002). The maximum likelihood parameter values were estimated using simulated annealing, and asymptotic 2-unit support intervals were used to assess the strength of evidence for individual parameter estimates (Goffe et al., 1994). The squared correlation coefficient (R^2) of the regression between observed and predicted values was used as a measure of the goodness-of-fit of each model. All analyses were performed

using R 3.0.3 software (R Development Core Team, 2014) and the ‘likelihood’ and ‘survival’ packages for R.

3. Results

3.1. General patterns of browsing-induced seedling damage and mortality

After 40 months of observation, browsing emerged as the most important cause of mortality for *B. pendula* – 40% of the seedlings of this species were browsed to death during the study period –, but mortality due to browsing never exceeded 15% for the three conifer species (Table 2). No significant differences in survival curves were observed between sites, i.e. browsing-induced mortality was similar at both elevations for each species (Fig. 2). Overall lethality (i.e., the proportion of browsed trees that ended up dying) was high (over 70%) but there were some differences across species, with the three conifer species (*A. alba*, *P. sylvestris* and *P. uncinata*) showing higher lethality values (average values ranging from 67% to 81%) than birch, the only study species with sprouting ability (Table 3).

3.2. Effect of mesohabitat on seasonal patterns of browsing-induced mortality

B. pendula seedlings were heavily browsed – both during winter and summer – at all the tested mesohabitats and comparable mortality rates were found across elevations and types of forest cover (Fig. 3). In contrast, the mortality rate due to browsing showed for *A. alba* and *P. uncinata* a marked seasonal effect, with an interaction between season and elevation. During winter, browsing-induced mortality for these two species was observed on both elevational stages but only on seedlings planted in gaps, but the trend was the opposite during summer, when a higher probability of being browsed was observed for understory seedlings although only in the subalpine stage (Fig. 3). Finally, browsing rates on *P. sylvestris* were rather low, and did not result in significant higher mortality rates at any environmental condition.

3.3. Effect of microhabitat on browsing-induced mortality

The distance to the nearest shrub exerted a positive effect on the probability of mortality due to browsing of *P. uncinata* and

Table 2

Observed mortality rates by apparent cause for seedlings of four species of trees planted at montane and subalpine elevational belt.

	Browse-induced mortality		Lack of adaptation		Random causes	
	Montane	Subalpine	Montane	Subalpine	Montane	Subalpine
<i>Abies alba</i>	0.07	0.13	0.25	0.11	0.03	0.15
<i>Betula pendula</i>	0.42	0.48	0.15	0.22	0.03	0.17
<i>Pinus sylvestris</i>	0.08	0.09	0.24	0.40	0.04	0.15
<i>Pinus uncinata</i>	0.09	0.07	0.15	0.13	0.02	0.20




Fig. 2. Survival curves for seedlings of the four tree species planted at the two experimental sites over the course of the study period, based on Kaplan–Meier estimates of browsing-induced mortality. Black and gray lines represent seedlings in the montane and subalpine belts, respectively. Solid lines are Kaplan–Meier estimates whereas dashed lines are 95% confidence intervals. *P*-values indicate significance of the log-rank test between sites for each species. Shaded areas in the x-axis correspond to the growing season (from May to October). Note that the y-axis starts at 0.4 for greater clarity and easier comparison among species.

Table 3

Observed browsing rate (percentage of browsed plants relative to the total number of unprotected seedlings) and lethality (percentage of browsed seedlings that died within the next months) for seedlings of four species of trees planted at montane and subalpine elevational belt, and average for both elevations.

	Montane belt		Subalpine belt		Average	
	% Browsed seedlings	% Lethality	% Browsed seedlings	% Lethality	% Browsed seedlings	% Lethality
<i>Abies alba</i>	12.5	77.8	23.6	70.6	18.1	73.1
<i>Betula pendula</i>	65.0	55.6	63.9	56.5	69.4	56.0
<i>Pinus sylvestris</i>	11.1	62.5	18.1	69.2	14.5	66.7
<i>Pinus uncinata</i>	9.8	85.7	12.5	77.8	11.1	81.2

A. alba plants (Fig. 4), meaning that a seedling planted further from a shrub had greater chances of being browsed. For *A. alba*, this effect was modulated by elevation, and the response was higher at the subalpine belt, where the modeled probability of mortality increased from almost zero for a seedling growing at less than 50 cm from a shrub, to almost 1 for seedlings at more than 2.5 m apart from a shrub (Fig. 4). For *Betula* and *P. sylvestris*, the model that included distance to shrubs did not have stronger empirical support than the null model (Table 4), so an effect of distance to shrub on mortality for these species was discarded. This was also the case when the explanatory variables were herb abundance or light availability, and an effect of these variables on mortality could not be found for any of the species (data not shown).

4. Discussion

4.1. Effects of micro- and meso-habitat on browsing patterns

We found both micro- and meso-habitat factors to exert a significant effect on browsing patterns and associated seedling mortality. At the microhabitat level we found, as expected, the browsing-induced mortality of *P. uncinata* and *A. alba* to decrease when seedlings were growing close to a shrub. The role of shrubs as facilitators of tree regeneration has been widely observed in mountain areas, where shrubs act as ‘biotic refuges’ for seedlings (Heuze et al., 2005; Milchunas, 2002; Vandenberghe, 2007). In Mediterranean mountains, habitat amelioration has been pointed out as the main mechanism behind this facilitative effect (Castro




Fig. 3. Seasonal browsing-induced mortality (mean \pm SE) for seedlings of *Abies alba*, *Betula pendula*, *Pinus sylvestris* and *Pinus uncinata* as a function of elevational belt (montane vs. subalpine) and type of cover (forest cover vs. gap). Asterisks above bars indicate values of mortality significant different from zero, whereas values above bars indicate the significance of the difference between mortality of seedlings growing under forest cover and gaps for a given species and elevation * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$. Values correspond to the mean at the end of the winter (month 32) and summer (month 39) periods.

et al., 2004; Gómez-Aparicio and Canham, 2008). This was also observed in a previous work conducted in the same area, in which we found a protective role of shrubs on seedlings face to adverse climatic events (Ameztegui and Coll, 2013). The results of the present study indicate that, in addition to habitat amelioration, shrubs also increase plant survival by providing physical protection from herbivores, denoting that both mechanisms co-occur in Mediterranean mountain systems (as also found by Boulant et al., 2008). Contrary to some recent works conducted in other mountain systems (e.g. Häslér and Senn, 2012; Kupferschmid et al., 2013), we did not find grass abundance to affect browsing probability in our study area. The high abundance of grasses present in all our plots is probably at the origin of this divergence.

Contrary to our hypotheses, the patterns of seedling consumption differed between summer and winter at the mesohabitat scale. During summer, *Betula* seedlings were heavily browsed in all conditions, whereas *P. uncinata* and *A. alba* were mostly browsed when growing in the forest understory. These areas are frequently visited by livestock, which commonly seek shade in the warmest hours of the day. They also have low shrub densities (especially in the sub-alpine belt), which results in a greater visibility of the plants and hence in a greater chance of being browsed. During winter, the opposite consumption pattern was found, and mortality associated to browsing was in general higher in plots located in gaps. The different duration of snow cover in the plots established in these areas might explain the observed patterns. Faster melting snow-pack in gaps due to higher insolation can cause the seedlings to become available earlier in the year, during a period with potential scarcity of alternative food sources (Häslér and Senn, 2012; Jackson, 2009), hence increasing the pressure on them.

4.2. Impact of browsing on species-specific seedling mortality and forest dynamics

Browse-induced mortality rate during winter (i.e. caused exclusively by ungulates) was similar to browse-induced mortality observed during summer, when plots can be frequented by both wild ungulates and domestic livestock. However, intra-guild competition was probably occurring in the study area during summer, when the presence of cattle forced the ungulates to take refuge in the rocky slopes located above the main grazing areas. Thus, assuming that most of the summer browsing-induced mortality is caused by livestock, our results indicate that livestock exert a lower browsing pressure than wild ungulates at current grazing conditions, as hypothesized. Livestock are usually characterized as 'grazers', since they tend to avoid the consumption of woody plants except in situations with very high stocking densities or fodder scarcity (Mayer, 2006; Vandenbergh, 2007). The high overall herb availability in the study area during summer may thus have contributed to the moderate browsing pressure observed for livestock. On the other hand, wild ungulates are more selective and cause more abundant and severe damages than cattle to seedlings, often being described as 'browsers' (Didion et al., 2009; Liss, 1988; Mayer, 2005b, 2006). The browsing behavior is particularly important during winter, when other food resources are scarce and seedlings constitute a significant part of their diet (Häslér and Senn, 2012; Jackson, 2009; Motta, 1996).

Herbivores showed a clear preference for birch seedlings throughout the whole study period, both during winter and summer. Mortality values for *Betula* seedlings were much higher than expected, since this species is not commonly considered as one of the preferred options of browsers (see, for example, the reviews in Gill (1992) or Didion et al. (2011)). The higher size of birch plants compared to the other species could in part explain the observed differences, since tall plants are browsed more often than smaller ones, the latter being often obscured by surrounding vegetation or just ignored due to their small energy return (Kupferschmid et al., 2013; Senn and Suter, 2003). In our study area, a positive relationship between seedling height and the probability of being browsed was detected for our dataset when all species were analyzed together (data not shown), but we could not detect this effect for any of the species when analyzed separately, due to the small intraspecific variability in plant height.

For the three conifer species, browsing-induced mortality rates observed in the study area were low or moderate, as they never exceeded 15%. The low browsing pressure observed on both pines matches previous studies that suggest a low palatability of these species (Didion et al., 2011). In contrast, fir is considered one of the preferred species by ungulates, which cause significant




Fig. 4. Predicted variation in mortality risk for seedlings of *Abies alba*, *Betula pendula*, *Pinus sylvestris* and *Pinus uncinata* as a function of the distance to the nearest shrub. Black and gray lines represent seedlings in the montane and subalpine belts, respectively. Solid lines correspond to the predicted model, whereas dashed lines represent the interval defined by the parameter estimates ± 2 likelihood units. Horizontal lines indicate a lack of effect of the explanatory variable, and are only shown for comparative purposes.

Table 4

Summary of the models predicting browsing-induced mortality as a function of micro-habitat conditions for seedlings of four species of trees planted at two experimental sites in the Eastern Pyrenees. For each species and explanatory variable, AIC of the best-fit model is provided for all seedlings pooled together (AIC_p) and separated into two groups (montane vs. subalpine, AIC_s).

Variable	Model	AIC_p	AIC_s	R^2
<i>Abies alba</i>				
Dist. to shrubs	Lin	78.06	73.91	0.39
Null	Null	89.85	83.33	0.13
<i>Betula pendula</i>				
Null	Null	52.05	52.51	0.02
<i>Pinus sylvestris</i>				
Null	Null	59.16	57.76	0.02
<i>Pinus uncinata</i>				
Dist. to shrubs	Power	52.19	61.10	0.17
Null	Null	63.37	63.24	0.03

For each species, only models with stronger empirical support than the null model (i.e. an AIC_p at least two units lower) are provided. Lower AIC_s values than AIC_p values indicate stronger empirical support for separated data than for pooled data, i.e. an elevational effect. Models are: Lin = linear; Exp = exponential, Pow = power. Details on the model formulation, parameter estimates and 2-units support intervals for the best-fit models are provided in [Appendix A](#).

difficulties for its regeneration in mountain areas of Central Europe (Ammer, 1996; Heuze et al., 2005; Motta, 1996; Senn and Suter, 2003). In this study we unexpectedly found low browsing rates on fir seedlings, but this could probably be explained by the small size reached by the plants four years after planting.

For *A. alba*, *P. sylvestris* and *P. uncinata*, browsing induced similar or lower mortality than that observed for other causes, such as lack of climatic adaptation or sensitivity to extreme climatic events (Ameztegui and Coll, 2013). Therefore, one might conclude that

browsing does not play a decisive role in the dynamics of these species, or at least not to the point of limiting or preventing their regeneration. For birch, which concentrated most of the damage by browsing, herbivory could be more limiting, especially considering the pioneer nature of this species, which normally regenerates abundantly in forest gaps after disturbances.

Although the mortality for the three conifers was not high, the lethality (i.e. the ratio of dead trees to damaged trees) was very high for all the species, and most seedlings died within a few months after being browsed. High lethality values associated to browsing are common at early life-stages (Senn and Suter, 2003), contrary to what occurs at more advanced stages, when mortality associated to browsing only happens in the most extreme cases (Häsler et al., 2008; Kupferschmid and Bugmann, 2013). In this study, 2-year old seedlings were planted at the limits of their distribution, and therefore at the limits of their physiological tolerance, as is apparent from their low growth rate and high mortality after adverse climatic events (Ameztegui and Coll, 2013). This may explain in part the high lethality we found given that the ability of plants to get over browsing is known to strongly depend on tree vigor (Kupferschmid et al., 2013, 2014).

4.3. Conclusions

The predation rate observed in this study was overall low or moderate, and should not represent a constraint for the regeneration of the study species. However, it is worth noting that this study focused on the first years of plant development, when plants are still relatively small. At higher plant developmental stages, more important browsing damages may occur, particularly if ungulate populations continue to grow favored by land-use changes.

Our results stressed important species-specific, seasonal, and habitat-related differences in plant damage by herbivores, which when occurred, was particularly lethal. The information derived from this study can be used by forest managers to adapt their practices and reduce browsing damages. For example, at the micro-habitat scale we showed that shrubs play an important role protecting seedlings from browsing. The known ‘nursing’ effect of shrubs has already been suggested as a technique to increase the success of reforestation plans in Mediterranean mountains (Castro et al., 2002; Gómez-Aparicio et al., 2008), and could also be used to promote species diversification of pine forests. Managers can control the understory shrubby vegetation through an adequate management of the canopy cover (Coll et al., 2010), seeking the conditions that minimize damages by herbivores. In a future context of increasing browsing pressure by wild herbivores, further knowledge in this topic may help foresters to define adequate forest management and planning strategies to allow regeneration success in these areas. Further studies might also be required to elucidate if the habitat-related consumption patterns observed in our study area are consistent across different forest types and browsing pressure gradients.

Acknowledgements

This research was primarily supported by the Spanish Ministry of Science and Innovation via the projects DINAMIX (AGL2009-13270-C02) and RESILFOR (AGL2012-40039-C02-01) and by the European Commission through the Marie Curie IRSES project “NEWFORESTS”. The Spanish Ministry of Science provided LC with support through a Ramon y Cajal contract (RYC-2009-04985), and the Spanish Ministry of Education provided AA with support through a predoctoral grant (FPU Program – AP2007-01663). The authors are particularly grateful to A. Bargañés, S. Martín, L. Ivorra and M. Pallarés for their invaluable work during field sampling and laboratory processing. F. Cano helped us to find the most suitable forests for this study, and the Socarrel team kindly offered their facilities during the fieldwork stage. We also thank the ‘Parc Natural del Cadí-Moixeró’ and the ‘Montellà i Martinet’ municipality for kindly giving permission to access the park and sample the data.

Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.foreco.2015.01.021>.

References

- Alía, R., Alba, N., Agúndez, D., Iglesias, S., 2005. Manual para la comercialización y producción de semillas y plantas forestales. Materiales de base y de reproducción. Serie Forestal. DGB, Madrid, Spain.
- Ameztegui, A., Coll, L., 2013. Unraveling the role of light and biotic interactions on seedling performance of four Pyrenean species under environmental gradients. *For. Ecol. Manage.* 303, 25–34.
- Ameztegui, A., Brotons, L., Coll, L., 2010. Land-use changes as major drivers of mountain pine (*Pinus uncinata* Ram.) expansion in the Pyrenees. *Glob. Ecol. Biogeogr.* 19, 632–641.
- Ammer, C., 1996. Impact of ungulates on structure and dynamics of natural regeneration of mixed mountain forests in the Bavarian Alps. *For. Ecol. Manage.* 88, 43–53.
- Apollonio, M., Andersen, R., Putman, R. (Eds.), 2010. European Ungulates and their Management in the 21st Century. Cambridge University Press, Cambridge, UK.
- Balcells, R., 1983. Evolución socio-económica reciente de tres comunidades comarcas pirenaicas y destino actual de las superficies más productivas de su demarcación. *Cuad. Invest. Geográfica* 9, 41–82.
- Bergström, R., Edenius, L., 2003. From twigs to landscapes – methods for studying ecological effects of forest ungulates. *J. Nat. Conserv.* 10, 203–211.
- Boulant, N., Navas, M.-L., Corcket, E., Lepart, J., 2008. Habitat amelioration and associational defence as main facilitative mechanisms in Mediterranean grasslands grazed by domestic livestock. *Ecoscience* 15, 407–415.
- Bugmann, H., Weisberg, P.J., 2003. Forest–ungulate interactions: monitoring, modeling and management. *J. Nat. Conserv.* 10, 193–201.
- Burnham, K.P., Anderson, D., 2002. Model Selection and Multi-Model Inference: A Practical Information-Theoretic Approach. Springer-Verlag, New York, USA.
- Casals, P., Baiges, T., Bota, G., Chocarro, C., de Bello, F., Fanlo, R., Sebastià, M.T., Tauli, M., 2009. Silvopastoral systems in the Northeastern Iberian Peninsula: A multifunctional perspective. In: Rigueiro-Rodríguez, A., McAdam, J., Mosquera-Losada, M. (Eds.), *Agroforestry in Europe SE - 8, Advances in Agroforestry*, Springer Netherlands, pp. 161–181.
- Castro, J., Zamora, R., Hódar, J.A., Gómez, J.M., 2002. Use of shrubs as nurse plants: a new technique for reforestation in Mediterranean mountains. *Restor. Ecol.* 10, 297–305.
- Castro, J., Zamora, R., Hódar, J.A., Gómez, J.M., Gómez-Aparicio, L., 2004. Benefits of using shrubs as nurse plants for reforestation in Mediterranean mountains: a 4-year study. *Restor. Ecol.* 12, 352–358.
- Coll, L., González-Olabarria, J.R., Mola-Yudego, B., Pukkala, T., Messier, C., 2010. Predicting understory maximum shrubs cover using altitude and overstory basal area in different Mediterranean forests. *Eur. J. For. Res.*
- Danell, K., 2003. Ungulates as drivers of tree population dynamics at module and genet levels. *For. Ecol. Manage.* 181, 67–76.
- DGMNB, 2014. *Memòria Anual de Caça Temporada 2012–2013*, Barcelona, Spain.
- Didion, M., Kupferschmid, A.D., Bugmann, H., 2009. Long-term effects of ungulate browsing on forest composition and structure. *For. Ecol. Manage.* 258, S44–S55.
- Didion, M., Kupferschmid, A.D., Wolf, A., Bugmann, H., 2011. Ungulate herbivory modifies the effects of climate change on mountain forests. *Clim. Change* 109, 647–669.
- García-Ruiz, J.M., Lasanta, T., 1990. Land-use changes in the Spanish Pyrenees. *Mt. Res. Dev.* 10, 267–279.
- Gill, R.M.A., 1992. A review of damage by mammals in northern temperate forests. 1. Deer. *Forestry* 65, 145–169.
- Goffe, W.L., Ferrier, G.D., Rogers, J., Biersens, H., Booty, M., Goldberg, D., Kerr, S., Levy, D., Maasoumi, E., Mccaffrey, D., 1994. Global optimization of statistical functions with simulated annealing. *J. Econom.* 60, 65–99.
- Gómez-Aparicio, L., Canham, C.D., 2008. Neighborhood models of the effects of invasive tree species on ecosystem processes. *Ecol. Monogr.* 78, 69–86.
- Gómez-Aparicio, L., Zamora, R., Castro, J., Hódar, J.A., 2008. Facilitation of tree saplings by nurse plants: microhabitat amelioration or protection against herbivores? *J. Veg. Sci.* 19, 161–172.
- Häsler, H., Senn, J., 2012. Ungulate browsing on European silver fir *Abies alba*: the role of occasions, food shortage and diet preferences. *Wildlife Biol.* 18, 67–74.
- Häsler, H., Senn, J., Edwards, J., 2008. Light-dependent growth responses of young *Abies alba* to simulated ungulate browsing. *Funct. Ecol.* 22, 48–57.
- Heuze, P., Schnitzler, A., Klein, F., 2005. Is browsing the major factor of silver fir decline in the Vosges mountains of France? *For. Ecol. Manage.* 217, 219–228.
- Jackson, J., 2009. The annual diet of the Roe deer (*Capreolus capreolus*) in the New Forest, Hampshire, as determined by rumen content analysis. *J. Zool.* 192, 71–83.
- Körner, C., Paulsen, J., 2004. A world-wide study of high altitude treeline temperatures. *J. Biogeogr.* 31, 713–732.
- Kräuchi, N., Brang, P., Schönenberger, W., 2000. Forests of mountainous regions: gaps in knowledge and research needs. *For. Ecol. Manage.* 132, 73–82.
- Kupferschmid, A.D., Bugmann, H., 2013. Timing, light availability and vigour determine the response of *Abies alba* saplings to leader shoot browsing. *Eur. J. For. Res.* 132, 47–60.
- Kupferschmid, A.D., Zimmermann, S., Bugmann, H., 2013. Browsing regime and growth response of naturally regenerated *Abies alba* saplings along light gradients. *For. Ecol. Manage.* 310, 393–404.
- Kupferschmid, A.D., Wasem, U., Bugmann, H., 2014. Light availability and ungulate browsing determine growth, height and mortality of *Abies alba* saplings. *For. Ecol. Manage.* 318, 359–369.
- Lasanta, T., 1990. Tendances actuelles de l’organisation spatiale des montagnes espagnoles. *Ann. Geogr.* 551, 51–71.
- Liss, B.M., 1988. Der Einfluss von Weidevieh und Wild auf die natürliche und künstliche Verjüngung im Bergmischwald der ostbayerischen Alpen. *Forstwissenschaftliches Cent* 107, 14–25.
- Mayer, A.C., 2005a. Experimental determination of the effects of cattle stocking density and grazing period on forest regeneration on a subalpine wood pasture. *Anim. Res.* 54, 153–171.
- Mayer, A.C., 2005b. Long-term impact of cattle grazing on subalpine forest development and efficiency of snow avalanche protection. *Arct. Antarct. Alp. Res.* 37, 521–526.
- Mayer, A.C., 2006. Influence of cattle stocking rate on browsing of Norway spruce in subalpine wood pastures. *Agrofor. Syst.* 66, 143–149.
- Mayer, A.C., Stöckli, V., Huovinen, C., Konold, W., Estermann, B.L., Kreuzer, M., 2003. Herbage selection by cattle on sub-alpine wood pastures. *For. Ecol. Manage.* 181, 39–50.
- Milchunas, D.G., 2002. Grazing refuges, external avoidance of herbivory and plant diversity. *Oikos* 99, 113–130.
- Motta, R., 1996. Impact of wild ungulates on forest regeneration and tree composition of mountain forests in the Western Italian Alps. *For. Ecol. Manage.* 88, 93–98.

- Navarro-González, I., Pérez-Luque, A.J., Bonet, F.J., Zamora, R., 2013. The weight of the past: land-use legacies and recolonization of pine plantations by oak trees. *Ecol. Appl.* 23, 1267–1276.
- Parent, S., Messier, C., 1996. A simple and efficient method to estimate microsite light availability under a forest canopy. *Can. J. For. Res.* 26, 151–154.
- Poyatos, R., Latron, J., Lloret, F., 2003. Land use and land cover change after agricultural abandonment – the case of a Mediterranean mountain area (Catalan Pre-Pyrenees). *Mt. Res. Dev.* 23, 362–368.
- Putman, R.J., 1996. Ungulates in temperate forest ecosystems: perspectives and recommendations for future research. *For. Ecol. Manage.* 88, 205–214.
- Putman, R., Apollonio, M., Andersen, R. (Eds.), 2011. *Ungulate Management in Europe: Problems and Practices*. Cambridge University Press.
- R Development Core Team, 2014. *R: A language and environment for statistical computing*.
- Reimoser, F., 1996. Impact of ungulates on forest vegetation and its dependence on the silvicultural system. *For. Ecol. Manage.* 88, 107–119.
- Reimoser, F., 2003. Steering the impacts of ungulates on temperate forests. *J. Nat. Conserv.* 252, 243–252.
- Senn, J., Suter, W., 2003. Ungulate browsing on silver fir (*Abies alba*) in the Swiss Alps: beliefs in search of supporting data. *For. Ecol. Manage.* 181, 151–164.
- Vandenberghe, C., 2007. Short-term effects of cattle browsing on tree sapling growth in mountain wooded pastures. *Plant Ecol.* 188, 253–264.
- Weisberg, P.J., Bugmann, H., 2003. Forest dynamics and ungulate herbivory: from leaf to landscape. *For. Ecol. Manage.* 181, 1–12.