



Analyzing the recent dynamics of wildland fires in *Quercus suber* L. woodlands in Sardinia (Italy), Corsica (France) and Catalonia (Spain)

Michele Salis¹ · Bachisio Arca¹ · Fermin Alcasena-Urdiroz² · Antonella Massaiu³ · Valentina Bacciu⁴ · Frédéric Bosseur⁵ · Philippe Caramelle³ · Sandro Dettori⁶ · Ana Sofia Fernandes de Oliveira⁶ · Domingo Molina-Terren⁷ · Grazia Pellizzaro¹ · Paul-Antoine Santoni⁵ · Donatella Spano^{4,6} · Cristina Vega-Garcia^{2,8} · Pierpaolo Duce¹

Received: 17 July 2018 / Revised: 31 January 2019 / Accepted: 6 February 2019 / Published online: 20 February 2019
© The Author(s) 2019

Abstract

Wildland fires represent a major threat to *Quercus suber* L. ecosystems, which provide relevant socioeconomic and ecological services in the Mediterranean Basin. In this work, we analyzed recent wildland fire dynamics in cork oak woodlands along the fire-prone areas of Sardinia (Italy), Corsica (France) and Catalonia (Spain). We first characterized geographic extent and main characteristics of cork oak woodlands in these regions and analyzed how environmental (climate and elevation) and socioeconomic factors (population and land uses) vary in the areas covered by *Quercus suber* L. We then evaluated how wildfires affected cork oak stands and, by logistic regression analysis, to what extent wildfires in cork oak areas were related to the above set of environmental and anthropic explanatory variables. Results revealed specific variations across study areas in cork oak characteristics as well as in environmental and social factors. We highlighted the spatial and temporal patterns of wildfires on cork oak woodlands, in terms of extent, seasonality, frequency and main driving factors. In the period 2003–2015, the percentage of cork oak woodlands burned ranged from 3.42% in Corsica to 11.30% in Sardinia. A few large and severe wildfires accounted for most of the area burned in cork oak woodlands. The most significant predictive variable that explained the spatial variation in wildland fire ignitions inside or nearby cork oaks was summer precipitation, while the weight of other factors varied depending on the region. This study provides evidence on recent fire dynamics in cork oak woodlands and gives valuable information and insights for the implementation of forest management and planning strategies in the Mediterranean area.

Keywords Cork oaks · Mediterranean Basin · Fire regime · Fire management · Land management

Introduction

Cork oak (*Quercus suber* L.) is a long-lived (up to hundreds of years) evergreen tree, endemic to the Mediterranean basin (EEA 2007). Cork oak woodlands are mostly concentrated in a few Mediterranean countries and cover about 2.2 million hectares (FAO 2013). The area of *Quercus suber* L. in

Europe is estimated to be close to 1.5 million hectares, the majority of which located in the Iberian Peninsula (FAO 2013; Dettori and Filigheddu 2016). The high presence of *Quercus suber* L. in the Mediterranean Basin is linked to its high economic and cultural value for rural communities (Urbietta et al. 2008; FAO 2013; Corona et al. 2018): cork is the second most important marketable non-wood forest product in the western Mediterranean Basin, with annual world cork market exports close to € 1.45 billion (Aronson et al. 2009; Catry et al. 2012a; APCOR 2016). *Quercus suber* L. stands have been intensively managed, protected and shaped by human activities over millennia for the nutritional values of acorns, to provide shelter and shade for livestock, and for the production of wood for domestic uses. In addition, cork oak forests supply a number of other goods and ecosystem services, including habitats for many

Communicated by Lluís Coll.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10342-019-01179-1>) contains supplementary material, which is available to authorized users.

✉ Michele Salis
miksalis@ibimet.cnr.it

Extended author information available on the last page of the article

animal species and plant endemisms, leisure activities, tourism, beekeeping, aromatic herbs or edible mushrooms. The social–ecological values that humans have imprinted on cork oak environments in the Mediterranean Basin generated unique cultural landscapes, which in turn influence local and regional culture and identity (Aronson et al. 2009; Plieninger et al. 2014). The landscape, cultural and ecological value of cork oak areas is also acknowledged in the Natura 2000 Network by the European Commission (DG Environment 2007).

As other Mediterranean forest systems, a number of previous studies have reported that cork oak ecosystems are in decline or endangered. The spatial distribution and vitality of *Quercus suber* L. woodlands are affected by several complex and sometimes interrelated disturbance factors (Luciano and Roversi 2001; Dettori and Filigheddu 2003, 2016; Silva and Catry 2006; EEA 2007; Pereira 2007; Acácio et al. 2009; Bugalho et al. 2011; FAO 2013; Oubrahim et al. 2015; Acha and Newing 2015; Oliveira et al. 2016; Dettori et al. 2018). Wildland fires represent one of the most relevant disturbances for cork oaks, as well as for other forests in the Mediterranean Basin. Nonetheless, it is widely accepted that fires are an integral part of the dynamics of many forest ecosystems, as they play a fundamental role in forest renewal and on insect and disease control (FAO 2013, Moritz et al. 2014). In Europe, forest fires mainly affect the Southern areas: in the period 2000–2015, about 51,000 fires per year have been recorded, burning approximately 380,000 ha of land (San-Miguel-Ayanz et al. 2016). In the Mediterranean Basin, most of the ignitions have anthropogenic origin, while area burned and fire intensity are primarily determined by weather conditions, topography, fuel load and continuity (Romero-Calcerrada et al. 2008; Pausas and Paula 2012; Ager et al. 2014b; Oliveira et al. 2014; Levin et al. 2016; San-Miguel-Ayanz et al. 2016; Turco et al. 2017; Rodrigues et al. 2018). Climate change is expected to increase future wildfire risk, area burned and the frequency of high severity events all over Southern Europe (Moriondo et al. 2006; Lung et al. 2013; Bedia et al. 2014; Kovats et al. 2014; Lozano et al. 2017). Furthermore, climate change can exacerbate stress conditions related to droughts, heat waves, pests and diseases, thereby contributing to the potential decline of cork oak woodlands in the next decades (Bergot et al. 2004; Luciano et al. 2005; Catry et al. 2012a; Costa et al. 2014). The recent trends of forest, agricultural and livestock activity abandonment, as well as the changes in the domestic energetic sources and the afforestation of abandoned lands, are responsible for the increase in highly flammable fuel accumulation and continuity, with potential effects on fire regime (Bonet and Pausas 2007; Baeza et al. 2011; Pausas and Fernandez-Munoz 2012; Madrigal et al. 2016). These factors could further increase the vulnerability to wildland fires of Mediterranean forests in future decades (Moreira et al. 2011; Salis et al. 2016b; Alcasena et al. 2016a).

Quercus suber L. is considered a fire-resistant and fire-resilient plant, due to its stem and crown resprouting capability (through epicormic buds), even after intense crown fires, and to the thick bark that ensures protection and insulation to the cambium from the heat produced by fires. Therefore, cork oaks are very competitive in the post-fire recovery phases in comparison with several other species (Pausas 1997; Curt and Pausas 2010; Catry et al. 2012a). For this reason, summer fires, even induced by landowners, have favored the expansion of cork oaks in Sardinia at the expense of *Quercus ilex* L. and *Quercus pubescens* Willd (Vogiatzakis and Careddu 2003). However, there is evidence that post-fire cork oak response and recovery capabilities may largely vary depending on several factors such as bark thickness, debarking interval, tree diameter, fire intensity and spread rate (Pausas 1997; Barberis et al. 2003; Moreira et al. 2007; Catry et al. 2009, 2012b). Moreover, an important issue for burned cork oaks is that the bark often becomes unsuitable for the most profitable industrial uses for a long period, which results in important economic losses to forest and rural communities. Hence, the incidence and effects of wildland fires on cork oak stands in the Mediterranean area are highly relevant to land managers, planners and policy-makers, according to the strategic economic, ecological and cultural importance of the *Quercus suber* L. woodlands.


In this work, we investigated the recent wildland fire dynamics on cork oak stands in three Mediterranean regions, namely Sardinia (Italy), Corsica (France), and Catalonia (Spain), where *Quercus suber* L. woodlands cover approximately 265,000 ha (about 18% of the European cork oak surface area). This research addresses the limited knowledge on recent wildfire trends in cork oak areas and on the geographic distribution of *Quercus suber* L. in the areas under investigation. The main research questions of this work are: (1) What are the main geographic, environmental and socioeconomic characteristics of the areas covered by *Quercus suber* L. woodlands? (2) To what extent have wildfires threatened cork oaks? (3) Do environmental and socioeconomic factors explain the historical wildfire incidence in *Quercus suber* L. woodlands? The answers to these questions provide insights and data on wildland fire dynamics that can support the implementation of forest management strategies and policies specific for cork oak forests.

Materials and methods

Study areas

The study was conducted in the fire-prone regions of Sardinia (Italy), Corsica (France) and Catalonia (Spain), located in the north-western part of the Mediterranean Basin (Fig. 1). For this study, we used “region” and “province”

Fig. 1 Digital elevation model (DEM) of the study areas (Catalonia, **a**; Sardinia and Corsica, **b**), along with the spatial distribution of *Quercus suber* L. woodlands (green polygons) and the administrative NUTS-3 boundaries (provinces of Cagliari, Sassari, Nuoro and Oristano in Sardinia; provinces of Barcelona, Tarragona, Girona and Lleida in Catalonia; departments of Corse du Sud and Haute-Corse in Corsica) used for the present study. The perimeters of the wildland fires (red color) that affected the study areas in the period 2003–2015 are also shown. (Color figure online)



as reference terms to identify NUTS-2 and NUTS-3 levels (EU Classification of Territorial Units for Statistics) for the study areas. The main characteristics of these areas are summarized in Table 1.

In the period 2003–2015, the total area burned in the study area was about 218,000 ha in Sardinia, 65,000 ha in Corsica and 61,000 ha in Catalonia; the average area burned per year was about 17,000 ha in Sardinia, 5,000 ha in Corsica and 4700 ha in Catalonia (Fig. 1 and Suppl. Fig. 1). The worst fire seasons in terms of total area burned (more than 30,000 ha) were observed in 2003 (Corsica), and in 2007 and 2009 (Sardinia). Overall, in the study period wildland fires affected less than 4% of the provincial areas in Catalonia as well as in Southern Corsica (Suppl. Fig. 1). Northern Corsica and Cagliari provinces were the most affected by fires (11.5% and 10.4% of the provincial areas in the 13-year study period, respectively). The highest provincial area burned by year was observed in Northern Corsica (6.1%) in 2003, Sassari (2.8%) in 2009 and Nuoro (2.3%) in 2007 (Suppl. Fig. 1). A peak in burned area was observed in July for all provinces, except for Tarragona (May), Northern Corsica and Barcelona (August) (Suppl. Fig. 2).

Data compilation

Cork oak woodland data

The *Quercus suber* L. woodlands data for the study areas were derived from the most accurate spatial input data available at national or regional levels. In Sardinia, the information source was the “Land Use Map of Sardinia” (2008),

which is a polygon shapefile that maps the Sardinian land uses with minimum mapping unit of 0.75 ha (www.sardegnageoportale.it). The information of this database is hierarchically organized following the CORINE Land Cover classification at the fifth level. The codes associated with cork oaks are: 2.4.1.3 (annual crops associated with permanent crops (pastures and grasslands with *Quercus suber* L. cover below 25%) and 3.1.1.2.2 (*Quercus suber* L. forests). The spatial location of cork oak woodlands in Corsica was derived from the “Inventaire Forestier National 2003” (<https://inventaire-forestier.ign.fr/>, minimum mapping unit of 20 ha). The map has three specific categories related to cork oak: *Quercus suber* L. forests, Mediterranean maquis with *Quercus suber* L., mixed *Quercus suber* L. woodlands and coppice. The cork oak information for Catalonia was obtained from the “Pure and Mixed Forest Type Map of Catalonia” (Vericat et al. 2010, minimum mapping unit of 25 ha). The classes associated with cork oaks, or with areas where cork oaks are the dominant forest types, are the following: a) *Quercus suber* L. forests; b) *Quercus suber* L. and other *Quercus* spp.; c) *Quercus suber* L. and *Pinus* spp.; d) *Quercus suber* L. and other species.

Wildland fire data

We gathered the perimeters of the wildland fires that affected Sardinia, Corsica and Catalonia for the study period 2003–2015. These data were provided by regional forest administrations: the Sardinia Forest Service (CFVA), the Office National de Forêt (ONF), and the Generalitat de Catalunya (GenCat), for Sardinia, Corsica

Table 1 Summary of the main characteristics of the study areas

	Sardinia (Italy)	Corsica (France)	Catalonia (Spain)
Size (km ²)	24,100	8700	32,100
Inhabitants	1.65 M	0.33 M	7.50 M
Population	Mostly concentrated nearby the main towns of Cagliari and Sassari	Mostly concentrated nearby the main towns of Ajaccio and Bastia	Mostly concentrated in the metropolitan area of Barcelona and the coastal areas
Climate type	Hot-summer Med.; warm-summer Med. above 750 m a.s.l.	Hot-summer Med.; warm-summer Med. above 500 m a.s.l.; cold-summer Med. above 1500 m. a.s.l.	Hot-summer Med.; warm-summer Med. in NW Catalonia, from cold-summer Med. until alpine above 1200 m a.s.l.
Annual precipitation	From ~400 mm in the southern coasts to ~1100 mm in the mountain areas	From ~500 mm in some coastal areas to more than 1100 mm in the mountain areas	From ~400 mm in the inner areas of Lleida to more than 1500 mm in the Pyrenees areas
Mean annual temperature	From ~7 °C in the mountain areas to ~17 °C along the southern coasts	From ~3 °C at the highest elevations to ~16 °C in the coastal areas	From ~0 °C in the Pyrenees to ~17 °C in the southern coasts

The sources of this information are provided in the next subsections

and Catalonia, respectively. We then merged and, in the case of unclosed polygons, rectified (about 0.01% of the data series) the wildland fire perimeters and the associated databases. Furthermore, we collected a longer time series (1990–2015) of wildland fire occurrence to analyze the spatial and temporal patterns of fire ignitions in the study areas. The ignition data of Catalonia were obtained from MAGRAMA (Spanish Ministry of Agriculture, Food and Environment), while for Sardinia and Corsica the data were provided by CFVA and ONF. We first checked the fire databases to evaluate their consistency, and we encountered a certain number of records (about 25%), particularly in the period 1990–2002, where ignition points coordinates were missing or incorrectly reported. In these cases, we randomly distributed the fire ignitions within each municipality of ignition. We then used the ignition point database to derive reference raster maps (100-m resolution) of historical ignition point densities for the periods 1990–2002 and 2003–2015. A radius search distance of 5000 m was set to generate smoothed maps of historical ignition density. This distance has been found significant in other works (i.e., Vega-Garcia et al. 1995) and compensates for possible errors in the ignition point coordinates of the historic databases.

Other input data

Terrain elevation information for the study areas was obtained from the EU-DEM data provided by the EEA (2017b). The EU-DEM is a 25-m resolution digital terrain model (DTM) and covers the whole EU countries plus some neighboring areas. Climate conditions of the study areas and of the cork oak stands were derived from the 1-km resolution “WorldClim version 2” database (Fick and Hijmans 2017), which provides monthly climate data for the period 1970–2000. Population density data at municipality level for the study areas were gathered from the Italian, French and Spanish census data and refer to the years 1990, 2000 (1999 for Corsica) and 2010. Finally, we used the Corine Land Cover data (CLC, years 1990, 2000, and 2012, EEA 2017a) to analyze temporal variations in land-use patterns for the study areas. The CLC data are seamless vector coverages of land covers for several European countries, with a minimum mapping unit of 25 ha. The CLC classification consists of an inventory of 44 land-cover classes, grouped in a three-level hierarchy. In this work, we focused our attention on the following classes: artificial surfaces (codes 111, 112, 121, 122, 123, 124, 131, 132, 133, 141, 142), arable land (211, 212), permanent crops (221, 222, 223), heterogeneous agricultural areas (241, 242, 243, 244), forests (311, 312, 313), scrub and/or herbaceous vegetation associations (321, 322, 323, 324).

Data analysis

Geographic extent and environmental and socioeconomic conditions in the cork oak areas

To characterize *Quercus suber* L. areas, we first mapped the cork oak woodlands and determined their total extent at provincial and regional levels while taking into account elevation classes and climate conditions. For terrain elevation, we used 250-m as reference range to discriminate the cork oak distribution in the study areas as a function of the elevation. We then applied the nonparametric Kruskal–Wallis statistical test to evaluate the significance of differences ($p=0.05$) among provinces. Regarding climatic conditions, we computed average, minimum and maximum values of temperatures and total precipitation at the provincial level and on the areas covered by cork oak woodlands, using as temporal reference: (1) the months from June to September (which essentially represent the fire season in the majority of the study areas) and (2) the annual averages. We then applied the Kruskal–Wallis test to evaluate the significance of differences ($p=0.05$) in climatic conditions among provinces. To evaluate demographic and land-use variations from 1990 to 2010 in the cork oak zones of the study areas, we selected municipalities characterized by a presence of *Quercus suber* L. woodlands above 5% of the total municipality area (Suppl. Fig. 3, and Figs. 2 and 3).

Historical trends in wildfires on cork oak areas

We intersected wildland fire perimeters and cork oak woodlands polygons to quantify the hectares of *Quercus suber*

L. burned at provincial and regional scales in the period 2003–2015. Moreover, we determined the cork oak area burned by year and by month in the study areas. The role played by the size of historical wildland fires (using as reference three size classes: ≤ 1 ha; 1–10 ha; 10–100 ha; > 100 ha) on cork oak area burned in the period 2003–2015, at the regional and provincial levels, was also investigated. We then evaluated the distance between ignition locations of large wildland fires and *Quercus suber* L. woodlands polygons by using the “Near” command of ArcGIS 10.3. For this purpose, we used 100 ha as a threshold to identify large events. The analysis was carried out considering two reference periods (2003–2015 and 1990–2002) and three reference distance classes (< 1 km, 1–10 km, and > 10 km).

Relationships between wildfire occurrence in cork oaks and environmental and socioeconomic factors

We applied logistic regression to analyze, at the regional scale, the relationships between the 1990–2015 wildland fires ignited inside or in the proximity of cork oak woodlands (using a 1-km buffer area around the *Quercus suber* L. polygons) and the following explanatory environmental and socioeconomic variables: climate, topography, population density and land use. We rasterized the explanatory variables using a reference grid of 5-km resolution for the whole study areas and quantified the number of annual fire ignitions for each cell. The logistic regression was used for binary classification (presence/absence) of fire ignitions inside the cork oak grid cells. The estimates of the binary response probability were based on the values of the explanatory variables. The final multivariate models were obtained by using a

Fig. 2 Map of the percent change in population between 1990 and 2010 in the study areas. The histograms report the number of inhabitants in the municipalities characterized by the presence of *Quercus suber* L. woodlands above 5% of the municipality area. The population data refer to census data of 1990, 2000, and 2010





Fig. 3 Map of the main land covers of the study areas according to Corine Land Cover 2012 (CLC2012). The histograms report the percent variations between CLC2012 and CLC1990 of the main land covers for the municipalities characterized by the presence of *Quercus suber* L. woodlands above 5% of the municipality area. A.S. = artificial surfaces; A.L. = arable land; H.A. = heterogeneous agricultural areas; P.C. = permanent crops; S.&H. = scrubs and herbs association; F. = forests; O. = other land covers



stepwise approach and considering all possible combinations of the variables for each region. The null hypothesis (the effects of the explanatory variables in the logistic models are equal to zero) was tested by the p value of the Wald Chi-square statistics. The Wald statistic was also used to evaluate the significance of both intercept and predictors used in the models. We evaluated the adequacy of the fitted models by the Hosmer and Lemeshow (2000) test, a Pearson Chi-square statistic that reflects the differences between observed and expected probabilities and provides an estimate of the lack of fit ($p=0.01$). In addition, the association between observed and predicted data was measured by the c -statistic (Bamber 1975), which is the percentage of pairs with concordant prediction and is commonly used as a standard measure of the predictive accuracy of a logistic regression model. C -statistic values were used to compare the different models, considering that c values range from 0.5 (random classification) to 1 (perfect association). We used the overall model accuracy to test the improvement determined by the use of explanatory variables over the intercept-only model.

Results

Characterizing geographic extent and environmental and socioeconomic conditions of the cork oak areas

As shown in Table 2 and Fig. 1, the presence of *Quercus suber* L. woodlands differs at the provincial level. Sardinia has about 138,400 ha of lands covered by cork oak (5.75% of the regional territory), while in Corsica the area covered by cork oak woodlands is around 64,500 ha (7.40% of the regional territory) (Table 2). On the other hand, *Quercus suber* L. woodlands encompass only about 1.91% (approximately 61,400 ha) of Catalonia (Table 2) and are only present in the northeastern zone of this region (Barcelona and Girona provinces).

In Sardinia, the northern province of Sassari accounts for about 60% of the Sardinian cork oak woodlands, while in Corsica *Quercus suber* L. woodlands are mainly

Table 2 Summary of the *Quercus suber* L. woodlands extension (in hectares) for the study areas. The relative percentage of provincial and regional area is reported under parenthesis

Cagliari	Sassari	Nuoro	Oristano	Sardinia
9030 (1.19%)	80,430 (10.46%)	35,880 (6.20%)	13,090 (4.33%)	138,430 (5.75%)
Corse du Sud	Haute-Corse			Corsica
46,570 (10.95%)	17,930 (4.01%)			64,500 (7.40%)
Barcelona	Girona	Tarragona	Lleida	Catalonia
7910 (1.02%)	53,480 (9.06%)	–	–	61,390 (1.91%)

The surface area is rounded to the nearest ten

present in Corse du Sud (46,570 ha) (Table 2). Catalonian *Quercus suber* L. woodlands are widespread in the province of Girona, which accounts to about 87% of the overall regional cork oak area (Table 2). Considering the distribution of cork oaks at the provincial level, Corse du Sud, Sassari and Girona show the highest presence of *Quercus suber* L. woodlands, which cover more than 9% of the provincial area. Cork oak woodlands account for 4–6% of the provincial territory in Nuoro, Oristano, and Haute-Corse, while their incidence in the other provinces is very limited (Cagliari and Barcelona) or null (Tarragona and Lleida) (Table 2).

Concerning the analysis of the cork oak stands at different elevation classes, the study areas show statistically significant differences (Table 3). *Quercus suber* L. woodlands are located at elevations much above the average provincial elevation in Cagliari and Oristano (376 m vs. 227 m a.s.l. and 350 m vs. 210 m a.s.l., respectively), while in Corsica and Catalonia they grow in areas below or largely below the average provincial elevation. In Corsica, more than 80% of *Quercus suber* L. woodlands is located in the 0–250 m

elevation class (Table 3). Also in Catalonia, the presence of cork oaks is most important at 0–250 m elevation class (about 60% of the total area), while only 33% of the Catalonia cork oak woodlands is found at 250–500 m altitude. The presence of cork oaks at elevations above 500 m is less than 5% of the regional and provincial areas of Catalonia and Corsica (Table 3). In Sardinia, about 50% of the *Quercus suber* L. stands are located at 250–500 m altitude class, with a relevant presence also at the 500–750 m elevation class (26% vs. less than 3% for Catalonia and Corsica). In the provinces of Nuoro and Cagliari, cork oak woodlands at elevations above 500 m represent about 55% and 40% of the provincial cork oak area, respectively. There are no significant statistical differences in average elevation of cork oak woodlands between Catalonian provinces, as well as for Cagliari, Oristano, and Sassari provinces (Table 3).

The average climatic conditions of *Quercus suber* L. woodlands at the provincial level are provided in Table 4. Overall, the climate of these areas is quite similar, particularly at the annual time lag (i.e., average temperatures range approximately from 14.9 °C in Oristano to 14.1 °C in

Table 3 Average terrain elevation (ELEV), average elevation of cork oak woodlands (ELEV QS) and percentage distribution of *Quercus suber* L. woodlands at different terrain elevation classes (0–250; 250–500; 500–750; > 750 m a.s.l.) in the provinces of the study areas

Province	ELEV (m)	ELEV QS (m)	0–250	250–500	500–750	> 750
Cagliari	227.1f	376.3b	24.08	36.54	38.84	0.54
Nuoro	526.1b	504.4a	12.32	32.57	41.37	13.74
Oristano	210.9g	350.0b	14.08	75.49	10.36	0.07
Sassari	325.7e	394.7b	22.39	55.33	20.00	2.28
Corse du Sud	560.4a	159.4e	87.51	12.13	0.36	0.00
Haute-Corse	592.3a	201.7d	69.07	28.30	2.48	0.15
Barcelona	518.1c	251.2c	68.78	29.14	2.07	0.00
Girona	517.2d	243.6c	60.08	34.05	5.16	0.71

The Kruskal–Wallis test ($p=0.05$) was performed to evaluate statistical differences in average elevation and average elevation of cork oak woodlands among provinces. Different letters in the same column indicate significant differences among provinces at $p=0.05$

Table 4 Average mean temperature (TAVG, in °C), minimum temperature (Tm, in °C), maximum temperature (TM, in °C), and accumulated precipitation (PP, in mm), at the annual level and for the period June–September, in the areas covered by *Quercus suber* L. woodlands

Province	Year				June–September			
	T	Tm	TM	PP	T	Tm	TM	PP
Cagliari	14.8b	10.6e	19.0b	624.2e	21.6b	16.5d	28.6b	76.4f
Nuoro	14.1e	10.3f	17.9g	716.4b	21.1d	16.4e	27.8c	90.4d
Oristano	14.9a	10.8d	19.1a	728.0b	21.8a	16.7c	28.9a	83.9e
Sassari	14.5c	10.8d	18.3d	703.5c	21.3c	16.7c	27.8c	88.1d
Corse du Sud	14.9a	11.8a	18.1e	612.7e	21.0e	17.4b	26.1f	91.8d
Haute-Corse	14.5d	11.0c	18.0f	663.6d	20.7f	16.7c	26.3e	113.1c
Barcelona	14.9ab	11.4b	18.4c	761.4a	21.6b	17.8a	26.9d	232.9a
Girona	14.7bc	11.0c	18.4cd	726.7b	21.3c	17.3b	26.9d	214.1b

The results are summarized at the provincial level. Climate data were obtained from the 1-km resolution “WorldClim version 2” data (Fick and Hijmans 2017). The Kruskal–Wallis test ($p=0.05$) was performed to evaluate statistical differences in climate conditions of cork oak woodlands among provinces. Different letters in the same column indicate significant differences at $p=0.05$

Nuoro). When taking into account only the June–September period, the ranges of some climatic variables in the *Quercus suber* L. woodlands show to be larger, particularly the average accumulated precipitation (i.e., from 76.4 mm in Cagliari to 232.9 mm in Barcelona) and the average maximum temperature (from 26.1 °C in Corse du Sud to 28.9 °C in Oristano) (Table 4). The maps of average maximum temperatures and accumulated precipitation observed in the study areas during the period June–September are presented in Suppl. Figs. 4 and 5.

The variations in population density in the municipalities where the presence of *Quercus suber* L. woodlands is above 5% of the municipality area are reported in Fig. 2, which is summarized at the regional level. Overall, the population in Corsica and Catalonia increased between 1990 and 2010 by 22% and 39%, respectively, with the highest variation observed in the province of Barcelona (+48%). On the contrary, in the same period we observed a decrease of about 5% of the inhabitants in the main cork oak areas of Sardinia, particularly in the inner lands. In addition, while in Catalonia and Corsica the percent change in population from 1990 to 2010 was quite uniform at the provincial level, Sardinia showed uneven changes: the province of Sassari exhibited slight changes and average variations close to zero, while the decrease in population in the main cork oak areas was over 11% in Cagliari and Oristano and close to 10% in Nuoro.

As for changes in main land uses between 1990 and 2012 in the municipalities characterized by the presence of *Quercus suber* L. woodlands above 5% of the municipality area (Fig. 3), we observed an overall increase in the surface areas covered by forests (particularly relevant in Catalonia) and artificial surfaces. Scrubs and herbaceous vegetation decreased in the study areas, with particular relevance in Catalonia and Sardinia. Concerning the other land uses, the variations were more complex and did not follow a common trend (Fig. 3).

Evaluating the wildfire trends in cork oak areas

In the study period 2003–2015, the total cork oak woodland area burned in the regions under investigation was around 23,500 ha (Fig. 1). Sardinia accounted for about 66% of the total cork oak area burned. In this island, on average about 1200 ha of *Quercus suber* L. stands (about 0.87% of the regional cork oak area) was burned each year (Fig. 4). In the 13-year study period, the total Sardinia cork oak area burned was about 15,500 ha. The worst seasons for *Quercus suber* L. woodlands were observed in 2007 and 2009, with about 2.79% (\approx 3850 ha) and 3.23% (\approx 4450 ha), respectively, of the total cork oak area burned. In Corsica (Fig. 4), only 3.42% (\approx 2200 ha) of the regional area covered by *Quercus suber* L. stands was burned in the study period. This means that, on average, the annual cork oak area burned




Fig. 4 *Quercus suber* L. woodlands area affected by wildland fires versus regional cork oak area for Catalonia, Corsica and Sardinia, for the study period 2003–2015

was about 165 ha, that is 0.26% of the total Corsica cork oak area. Furthermore, the annual percentage of *Quercus suber* L. woodlands burned over the total area covered by cork oaks was always less than 1%. In Catalonia, about 9.38% (\approx 5700 ha) of the area covered by *Quercus suber* L. woodlands was burned, with an annual average of 0.72% (about 450 ha) (Fig. 4). Nearly 7.5% (\approx 4400 ha) of the Catalonia *Quercus suber* L. woodland area was burned in 2012 and was almost entirely concentrated in the province of Girona (Fig. 4). Overall, the combination of wildland fire and cork oak woodland shapefiles in Sardinia, Corsica and Catalonia revealed that the *Quercus suber* L. areas which burned more than once in the period 2003–2015 were limited and ranged from 0 to less than 1% of the total cork oak area burned at the provincial level.


A small number (<2%) of large fire events (> 100 ha) accounted for most of the total cork oak area burned at the regional scale. The role played by large fires in *Quercus suber* L. woodlands burned was very significant in Catalonia (about 99% of the total cork oak burned determined by large events), while in Corsica and Sardinia they accounted for about 79% and 72%, respectively, of the total *Quercus suber* L. area burned (Table 5). This analysis highlighted some relevant differences also at the provincial level. For instance, the most relevant variation in the role played by the different fire size class was observed in Sardinian provinces. In fact, while in Sassari and Nuoro provinces the large fires (> 100 ha) accounted for more than 75% of the total cork oak area burned, in southern Sardinia these events were responsible for about 31% of the *Quercus suber* L. burned, and the highest contribution to cork oak area burned was due to the fire size class 10–100 ha (Table 5). Corsica also presented marked differences: large events accounted for about 88% of the total *Quercus suber* L. woodlands burned in Haute-Corse and for about 64% in Corse du Sud. In Catalonia, in both

Table 5 Percentage of *Quercus suber* L. woodlands burned by wildland fire size class (≤ 1 ha; 1–10 ha; 10–100 ha; > 100 ha), for the period 2003–2015, at the regional (in bold) and provincial levels

Province/Region	Percentage of QS burned by wildfire size class				Total
	≤ 1 ha	1–10 ha	10–100 ha	> 100 ha	
Cagliari	2.75	18.17	48.21	30.86	100.00
Nuoro	0.49	7.36	15.15	77.01	100.00
Oristano	0.78	14.56	20.80	63.85	100.00
Sassari	0.61	8.21	17.13	74.05	100.00
Sardinia	0.67	8.94	17.94	72.45	100.00
Corse du Sud	1.17	5.13	29.36	64.34	100.00
Haute-Corse	0.84	4.76	6.08	88.31	100.00
Corsica	0.97	4.91	15.41	78.71	100.00
Barcelona	0.00	0.00	0.00	100.00	100.00
Girona	0.00	0.00	1.32	98.68	100.00
Catalonia	0.00	0.00	1.28	98.72	100.00

Girona and Barcelona provinces, the fire size class > 100 ha caused more than 98% of the total cork oak area burned. A single forest fire event (Suppl. Fig. 6a) played a key role in determining the cork oak losses in Catalonia and particularly in Girona province. Specifically, on July 2012, the largest Catalonian wildland fire (La Jonquera, about 10,500 ha) of the period 2003–2015 burned approximately 4100 ha of *Quercus suber* L. woodlands, corresponding to about 94% of the total cork oak area burned in that year, and to about 72% of the total cork oak area burned in the region over the 13 years of reference. Another example of the significant role played by large events is represented by the wildland fire of Nuoro (Suppl. Fig. 6b) in Sardinia. This event (about 9000 ha), which was the largest Sardinian wildland fire of the summer season 2007, burned approximately 2600 ha of *Quercus suber* L. woodlands, which represent about 67% and 17% of the total cork oak area burned in 2007 and during the study period, respectively.

In Catalonia and Sardinia, *Quercus suber* L. woodlands were mostly affected by fires in July (more than 65% of the total cork oak area), followed by August (with about 13% and 19% of the total cork oak area burned in Corsica and Sardinia, respectively) (Fig. 5). The cork oak area burned in July was about 10,550 ha in Sardinia and 4200 ha in Catalonia (concentrated in the province of Girona). However, in Sardinia, relevant differences in the monthly distribution of the cork oak burned were observed: in the northern provinces of Sassari and Nuoro, about 90% of the cork oak stands burned in July and August, while in the other two provinces the wildfires occurred in these months accounted for about 75% of the total *Quercus suber* L. area burned. Also in Corsica, July was the month with the highest percentage (about 39%) of *Quercus suber* L. woodlands burned, but a relevant percentage (about 22%) of the cork oak woodlands

**Fig. 5** Percentage of the monthly distribution of the *Quercus suber* L. woodlands area affected by fires in Catalonia, Corsica and Sardinia, for the study period 2003–2015

burned was also observed in August and October (Fig. 5). The Corsican provinces presented large differences in the monthly distribution of the cork oak burned: in Corse du Sud, about 73% of the total cork oak area burned was concentrated in July, while in Haute-Corse the peak (about 87%) was observed in August–October.

The relation between the wildland fire ignition locations and the distance to cork oak stands was analyzed considering the reference period 2003–2015 and the previous 13-year timeframe (1990–2002). Figure 6 highlights the spatial variations in large fire event (> 100 ha) location and ignition point densities between the two periods. On the whole, the fire ignition point density dropped from 1990–2002 to 2003–2015 for most of the study areas, the main exception being south-western Sardinia (Fig. 6), which showed more areas with peaks above eight fire ignitions km^{-2} in the period 2003–2015. Moreover, large fire (> 100 ha) number and area burned decreased considerably in the study areas moving from the first to the second period (Fig. 6). For instance, the number of large fires dropped to 289 events (vs. 541 in the period 1990–2002) in Sardinia, to 47 (vs. 113) in Corsica, and to 75 (vs. 116) in Catalonia. The analysis of the percentage of fire ignition locations as function of the distance from cork oak woodlands and provinces for both 1990–2002 and 2003–2015 revealed that from the first to the second period, wildland fires ignited in the vicinity (< 1 km) of cork oak woodlands areas tended to decrease, except in the province of Girona (Suppl. Table 1). In the period 2003–2015, the likelihood of fire ignitions at the distance class 0–1 km from *Quercus suber* L. woodlands was over 40% of the total ignitions in the provinces of Girona, Nuoro, and Sassari, while Cagliari and Barcelona provinces showed values below 15%. The province of Barcelona presented the largest distances between fire ignitions and cork oak areas (Suppl. Table 1).




Fig. 6 Maps of the historic ignition point density (in $\#/km^2$) of the periods 1990–2002 (left) and 2003–2015 (right) for the study areas. The ignition point density was calculated using the Point Density tool

of ArcGIS (ESRI) and a radius search distance of 5,000 m. The black triangles indicate the ignition point location of the large wildland fire events (> 100 ha) that affected the study areas for the above periods

Analyzing the relationships between wildfire occurrence in cork oak areas and environmental and socioeconomic factors

The driving variables that mostly affected wildland fire ignitions inside and nearby (1-km buffer) cork oak areas at the regional scale were identified by a set of logistic regression models (Table 6). The Catalonia model was not statistically significant according to the Wald test, while the Sardinia

and Corsica models resulted highly statistically significant ($p=0.01$). However, the Corsica and Sardinia models exhibited a significant lack of fit according to the Hosmer & Lemeshow test (Table 6). Overall, the models showed values of correct classification between 55% and 73%, with the highest values observed in Corsica. *C*-statistic values range from 0.73 (Sardinia model) to 0.79 (Corsica and Catalonia models) and thus indicated an overall good predictive accuracy of the logistic models (Table 6). In the study areas, the most significant predictive variable that explained the

Table 6 Significance of the variables that affected wildland fire ignitions inside and nearby (1-km buffer) cork oak areas as provided by logistic regression models, considering the study period 1990–2015

		Sardinia	Corsica	Catalonia
Predictive variables	ELEV	−0.005 (190.7)***	0.03 (43.04)***	n.s.
	SLO	0.06 (10.32)**	n.s.	n.s.
	TMAXS	−1.22 (470.5)***	n.s.	n.s.
	TMS	n.s.	7.58 (43.04)***	n.s.
	PRECS	0.01 (5.65)***	3.62 (−0.03)***	−0.03 (3.00)***
	PREC	n.s.	n.s.	−0.11 (3.91)**
	POP	0.006 (83.41)***	−0.005 (9.95)***	n.s.
	LU	Agricultural −0.01*** Anthropic −0.33 (30.6)***	Agricultural −2.29*** Anthropic 4.11 (10.46)***	n.s.
	Model performance parameters	WCF	862***	122***
HLCF		35.95***	33.20***	16.91 n.s.
CORR		67.2	73.0	55.4
<i>c</i>		0.73	0.79	0.79

The values of the coefficient estimates, the Wald Chi-square values and the significance of the Chi-square statistics are reported for each predictive variable. Regarding land uses, the main land-use classes are reported if significant. ELEV=elevation; SLO=slope; TMAXS=average summer maximum temperatures; TMS=average summer mean temperatures; PRECS=average summer precipitation; PREC=average annual precipitation; POP=population density; LU=land uses. WCF=Wald test coefficient; HLCF=Hosmer & Lemeshow test coefficient; CORR=percentage of correct estimates; *c*=*c*-statistic value. ***= $p<0.01$; **= $p<0.05$; n.s.=not significant

spatial variation of wildland fire ignitions inside or nearby cork oaks was summer precipitation. In more detail, for both Sardinia and Corsica, the most significant predictive variables of the logistic models were land-use classes (anthropic and agricultural areas), population density, elevation and summer precipitation (Table 6). As far as climatic conditions are concerned, we also found a major effect of summer maximum temperatures in Sardinia and summer average temperatures in Corsica. Slope played a significant role for the Sardinia model. In Catalonia, the spatial variation of fire ignitions inside or nearby cork oak areas was mainly explained by climatic conditions, namely summer precipitation and annual precipitation (Table 6).

Discussion and conclusions

The main goal of this work was to analyze the recent wildland fire dynamics on cork oak areas in Sardinia (Italy), Corsica (France), and Catalonia (Spain). We aimed to provide answers to the following research questions: (1) What are the main geographic, environmental and socioeconomic characteristics of the areas covered by *Quercus suber* L. woodlands? (2) To what extent have wildfires affected cork oaks? (3) Do environmental and socioeconomic factors explain the historical wildfire incidence in *Quercus suber* L. woodlands?

The study areas encompass about 265,000 ha of *Quercus suber* L. woodlands, about 18% of the European cork oak surface area. Our study showed relevant differences in the spatial distribution of cork oaks in the regions under investigation. The presence of cork oak stands is mainly concentrated in the provincial areas of Corse du Sud, Sassari and Girona: these provinces account for more than 50% of the regional *Quercus suber* L. woodlands. Previous studies (Dettori and Filigheddu 2003; Pereira 2007; FAO 2013) have pointed out that the differences in cork oak extent within provinces and regions depend upon several interrelated factors, among which the most relevant are climatic conditions, income and jobs related to the cork industry, pressure of pastoral and agricultural activities, and urban sprawl. Our work showed that the climatic conditions in the cork oak woodlands of the study areas were relatively similar, but for summer precipitation and summer average maximum temperatures. This could explain the high presence of cork oak woodlands at elevation classes > 500 m a.s.l. in Sardinia, while in Corsica and Catalonia *Quercus suber* L. areas are largely concentrated at the lowest elevation classes (< 250 m a.s.l.). Differences in climatic conditions would also explain why Sardinian cork oaks are located at higher elevations than in Corsica and Catalonia. Drought conditions and the increasing extent and role played by agricultural expansion could further explain the limited presence of cork oaks in

southern Sardinia, or their absence in Tarragona and Lleida provinces.

The variation in population density between 1990 and 2010 assumed distinct patterns across regions. While at the regional level most Corsican and Catalan municipalities presented, save from a few localized inner areas, a general increase in population density, Sardinia showed a large decline of population density in municipalities far from coastlands and main towns. The population density decline observed in the cork oak areas of Sardinia can be related to a combination of socioeconomic issues, including unemployment, abandonment of rural and forest activities, ageing population, low birthrates, distance to main services and towns, and low incomes, which led to emigration of young people toward coastal areas and towns, or even outside, as observed in other Mediterranean areas (Romero-Calcerrada and Perry 2004; Figueiredo and Raschi 2011; Kerckhof et al. 2016; Forleo et al. 2017). For land-use variation from 1990 to 2012, we observed an increase in forests and artificial surfaces and a reduction in shrublands in the municipalities covered by large extensions of cork oak woodlands for Sardinia, Corsica and Catalonia. The highest increase in forest areas (about 15%) in the vicinity of cork oak polygons was observed in Catalonia, followed by Sardinia and Corsica (about 2–4%). These variations suggest that cork oaks or other forest types are gradually occupying areas previously covered by shrublands and marginal rangelands, with a potential increase in fuel load and continuity, particularly in the zones where population density is decreasing and vegetation succession is leading to scrub encroachment and new forest development (Arianoutsou 2001; Scozzafava and De Sanctis 2006; Moreira et al. 2011). In addition, the increased presence of artificial surfaces in the municipalities covered by large areas of cork oak woodlands indicates an expansion of urban interfaces in the wildlands.

We found that in the study areas approximately 23,500 ha of cork oak stands (about 9% of the total *Quercus suber* L. woodlands) were affected by wildland fires in the period 2003–2015. Sardinia accounted for about 66% of the cork oak woodland area burned. Evident interannual variability in the cork oak area burned was observed. Overall, the percentage of cork oak woodlands burned at least once ranged from 3.42% in Corsica to 11.30% in Sardinia for the study period 2003–2015. The highest cork oak area burned observed in Sardinia could be partially explained by the higher presence of herbaceous fuels and dehesas, which can favor the quick spread of large wildland fires under strong wind conditions (Salis et al. 2018). The role played by environmental factors on burned area in the Mediterranean Basin has been explored in previous works. There is evidence that a small number of large fires spreading with extreme weather conditions is responsible for the majority of the area burned during a fire season and that containment activities have minor

effectiveness on fire growth during these events (Pereira et al. 2005; Koutsias et al. 2012; San-Miguel-Ayanz et al. 2013; Alcasena et al. 2016b; Salis et al. 2016a). Large wildland fires often exhibit extreme behavior (i.e., high intensity, crown fire activity and spotting fire emission) and can produce damaging impacts on human life, assets and activities as well as severe disturbances to forest ecosystems. Our work confirms that wildland fires greater than 100 ha, which accounted for 1–2% of the total number of wildland fires, were associated with most of the area burned in *Quercus suber* L. woodlands in the study areas, except for southern Sardinia. For instance, in the study period, about 70% of the cork oak woodlands in Catalonia burned in a single fire event (La Jonquera, final size about 10,000 ha), and more than 90% of the La Jonquera fire area burned at high or very high severity (severicat.ctfc.cat). Similar findings were observed in Sardinia during the 2007 and 2009 fire seasons when extreme weather conditions and simultaneous wildland fires concentrated in a few days overwhelmed fire suppression capabilities and burned large portions of cork oak areas in the provinces of Nuoro and Sassari.

The present study also reveals that both the number of fire events and the area burned by wildland fires bigger than 100 ha, which are the primary drivers of fire regime in cork oak landscapes in the study areas, decreased considerably from 1990–2002 to 2003–2015, and that overall the distance between fire ignitions and cork oak woodlands areas increased. Furthermore, we observed an overall reduction in fire ignitions from 1990–2002 to 2003–2015 for almost the entire study areas, the main exception being in southern and western Sardinia. This is in line with previous studies that analyzed wildland fire trends in the Mediterranean Basin (Turco et al. 2016; Jiménez-Ruano et al. 2017; Curt and Frejaville 2018). The fact that large wildland fires impacting cork oak areas dropped in recent years can be explained by the improvements in fire monitoring and suppression capabilities, by the reduction in fire ignitions related to agropastoral causes and by a general shift of both people and fire ignitions toward coastal and main residential areas (Viedma et al. 2006; Chas-Amil et al. 2013; Salis et al. 2014; Curt and Frejaville 2018). Some studies emphasized that fire exclusion policies can substantially increase the vulnerability of many dry forests to uncharacteristically extensive and severe perturbations, particularly from wildfire and insects/disease, due to their emphasis on short-term outcomes (reducing annual area burned and fire ignitions) versus long-term goals (promoting more resilient landscapes and forests) (Piñol et al. 2005, 2007; Calkin et al. 2015; Stephens et al. 2016; Camia et al. 2017; Curt and Frejaville 2018; Thompson et al. 2018). These issues pose severe threats for fire management in future years in the study areas, since the reduction in forest management activities, in conjunction with the limited occurrence of small size and low severity fires in cork oak

areas, and the decrease in population in inner areas such as seen in Sardinia, could promote the accumulation of unmanaged high-load fuels, the expansion of ladder fuels and the increase in fuel connectivity. This can lead to more fire-prone landscapes in the Mediterranean area, to a potential increase in the likelihood of extreme behavior and severe wildland fire events, thus able to spread further and increase in size, and to more limitations for effective fire suppression of aerial and terrestrial forces (Moreno et al. 1998; Pausas and Vallejo 1999; Pausas and Fernandez-Munoz 2012; Salis et al. 2013; Ferreira-Leite et al. 2016). Therefore, it is likely that, in future years, wildland fires affecting forest areas will start increasingly far from the woodlands and will impact the forest stands after spreading for long distances, thus resulting in an overall increase in the so-called megafires, as dramatically observed in the European summer season 2017 (San-Miguel-Ayanz et al. 2013; Viegas et al. 2017). High-intensity wildland fires could ultimately trigger substantial changes on cork oak forest structures from old-growth forests with a commercial interest to post-fire shrubby coppices, which require long-time and expensive restoration efforts. Consequently, specific management strategies including fuel treatments should be planned and carried out by policymakers and land managers to reduce potential losses from future wildland fires in the Mediterranean Basin, in order to preserve important natural ecosystem heritages like the cork oak forests. A number of works highlighted that surface fuel reduction, silvicultural treatments, and better-planned land mosaics could significantly decrease fire risk in cork oak areas (Baeza et al. 2006; Santana et al. 2011; Catry et al. 2012b; Schaffhauser et al. 2012; Curt et al. 2013; Salis et al. 2016b, 2018). This brings to light the need for integrating large fire spread analysis and fire risk transmission with the design of appropriate wild-fire management strategies, particularly for protecting highly valued areas or habitats of endangered species (Ager et al. 2014a, 2017; Alcasena et al. 2017, 2018a; Thompson et al. 2017; Salis et al. 2016b, 2018). Prevention and protection strategies may also involve prescribed burning action and may allow the spread of low-intensity wildland fires when weather conditions are mild, in agreement with the proactive concept of integrated fire management (Molina et al. 2010; Fernandes et al. 2013; Montiel Molina and Galiana-Martín 2016; Alcasena et al. 2018b).

We observed that the highest cork oak area burned was concentrated during the months of July and August, which in Catalonia and Sardinia accounted for almost 80% of the total *Quercus suber* L. area affected by fires. Summer months are typically the hottest and driest of the year in the Mediterranean Basin (Flohn and Fantechi 1984). This confirms the key role of drought conditions, and particularly of dead and live fuel moisture, in the dynamics of fire regimes in Mediterranean Europe (Pellizzaro et al. 2007; Moreira et al.

2011; Turco et al. 2017). From this point of view, the logistic regression analysis performed in the study areas demonstrated that climatic factors, namely summer and annual precipitation, and maximum and average summer temperatures, were significantly related to wildland fire ignitions nearby cork oak woodlands. Furthermore, summer precipitation was a key driving factor in determining wildfires nearby cork oaks for the three study areas. The fact that wildfires occurring in dry years tend to burn larger areas than in wet years was also observed in other European fire-prone areas, as, for instance, in the Valencia region of Spain (Pausas 2004), in Portugal (Viegas and Viegas 1994) and in Greece (Dimi-trakopoulos et al. 2011; Xystrakis et al. 2014). Moreover, our study indicates that, while in Catalonia fire ignitions in the vicinity of cork oaks are mainly explained by summer and annual precipitation, Corsica and Sardinia showed more complex patterns: in these regions, elevation and land use play a significant role, together with population density and some climate variables. These results support the fact that, even if wildfire ignitions in Mediterranean areas are mostly determined by anthropic factors, a significant component of the variability in wildfire occurrence is explained by climatic conditions (Pausas 2004, Pereira et al. 2013). For these reasons, predicted climate change is likely to affect future fire dynamics in *Quercus suber* L. woodlands. With a higher frequency of extreme weather conditions, fire-prone landscapes can be increasingly vulnerable and able to sustain high-intensity and uncontrollable fire events, which can negatively affect cork oak post-fire recovery (Fernandes and Rego 1996; Baeza et al. 2006; Pausas and Paula 2012; Schaffhauser et al. 2012). In addition, under the predicted changes in climate conditions, cork oak ecosystems could be potentially replaced by more xeric species and pyrophytic shrublands and could shift northward or to higher elevations (Costa and Madeira 2011; Costa et al. 2014; Ibáñez et al. 2017; Varela 2017).

In conclusion, the cork oak woodlands of the study areas represent an important portion of the European *Quercus suber* L. ecosystems and a significant ecological, cultural and economic heritage for rural communities. In the study period 2003–2015, about 9% of the cork oak woodlands of the study areas have been affected by wildland fires. Though this represents a relatively small percentage of the total *Quercus suber* L. surface area, cork oaks burned can lead to relevant losses of ecosystem services and income, as well as to the death of monumental and ancient trees, which are key components of cultural landscapes and agro-pastoral areas in Sardinia, Corsica and Catalonia. Wildfires nearby cork oak forests were primary linked to the occurrence of dry summers, and the most of the *Quercus suber* L. area burned was related to a few large fire events associated with extreme weather conditions. It is thus likely that the future changes in climate will further increase wildfire risk in Mediterranean

forests, unless significant efforts are made to promote appropriate forest management and planning strategies. Socio-economic changes can even foster the occurrence of high-intensity wildland fires, due to the magnitude of land-use changes and the abandonment of forest and rural activities.

Acknowledgements This work was supported by the Sardinia Region under the project “Multifunzionalità delle foreste a quercia da sughero in Sardegna” (Multifunctionality of cork oak forests in Sardinia) funded within L.R. n. 7/2007 “Promozione della ricerca scientifica e della innovazione tecnologica in Sardegna/Tender no. 2”, and by the “Proterina-2” Project funded by the Italia Francia-Marittimo Programme. We would like to thank the Sardinia Forest Service, the Office Nationale de Foret, the Generalitat de Catalunya and the Area of Defense against Forest Fires (ADIF) of the Spanish Ministry of Agriculture, Food and the Environment (MAGRAMA) for providing fire data and information for Sardinia, Corsica and Catalonia. We also thank the anonymous reviewers whose comments helped improve this manuscript.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Acácio V, Holmgren M, Rego F, Moreira F, Mohren GM (2009) Are drought and wildfires turning Mediterranean cork oak forests into persistent shrublands? *Agrofor Syst* 76:389–400
- Acha A, Newing HS (2015) Cork oak landscapes, promised or compromised lands? A case study of a traditional cultural landscape in Southern Spain. *Human Ecol* 4(43):601–611
- Ager AA, Day M, Finney MA, Vance-Borland K, Vaillant NM (2014a) Analyzing the transmission of wildfire exposure on a fire-prone landscape in Oregon, USA. *For Ecol Manag* 334:377–390
- Ager AA, Preisler H, Arca B, Spano D, Salis M (2014b) Wildfire risk estimation in the Mediterranean area. *Environmetrics* 25(6):384–396
- Ager AA, Evers CR, Day MA, Preisler HK, Barros AMG, Nielsen-Pincus M (2017) Network analysis of wildfire transmission and implications for risk governance. *PLoS ONE* 12(3):e0172867
- Alcasena F, Salis M, Nauslar NJ, Aguinaga AE, Vega Garcia C (2016a) Quantifying economic losses from wildfires in black pine afforestations of northern Spain. *For Policy Econ* 73:153–167
- Alcasena FJ, Salis M, Vega-Garcia C (2016b) A fire modeling approach to assess wildfire exposure in central Navarra, Spain. *Eur J For Res* 135:87–107
- Alcasena FJ, Salis M, Ager AA, Castell R, Vega-García C (2017) Assessing wildland fire risk transmission to communities in Northern Spain. *Forests* 8(2):30. <https://doi.org/10.3390/f802030>
- Alcasena F, Ager AA, Bailey JD, Pineda N, Vega-Garcia C (2018a) Towards a comprehensive wildfire management strategy for Mediterranean areas: framework development and implementation in Catalonia, Spain. *J Environ Manag* 1:2–9. <https://doi.org/10.1016/j.jenvman.2018.10.027>

- Alcáscena FJ, Ager AA, Salis M, Day MA, Vega-García C (2018b) Optimizing prescribed fire allocation for managing wildfire risk in central Catalonia. *Sci Total Environ* 621:872–885
- APCOR (2016) APCOR's Cork Yearbook 2016. <http://www.apcor.pt/wp-content/uploads/2016/09/Boletim-estatistico-2016.pdf>
- Arianoutsou M (2001) Landscape changes in Mediterranean ecosystems of Greece: implications for fire and biodiversity issues. *J Mediterr Ecol* 2:165–178
- Aronson J, Pereira JS, Pausas JG (2009) Cork Oak Woodlands on the edge: ecology, adaptive management, and restoration. Island Press, Washington
- Baeza MJ, Raventós J, Escarré A, Vallejo VR (2006) Fire risk and vegetation structural dynamics in Mediterranean shrubland. *Plant Ecol* 187:189–201
- Baeza J, Santana VM, Pausas JG, Vallejo R (2011) Successional trends in standing dead biomass in Mediterranean basin species. *J Veg Sci* 22:467–474
- Bamber D (1975) The area above the ordinal dominance graph and the area below the receiver operating characteristic graph. *J Math Psychol* 12:387–415
- Barberis A, Dettori S, Filigheddu MR (2003) Management problems in Mediterranean cork oak forests: post-fire recovery. *J Arid Environ* 54:565–569
- Bedia J, Herrera S, Camia A, Moreno JM, Gutierrez JM (2014) Forest fire danger projections in the Mediterranean using ENSEMBLES regional climate change scenarios. *Clim Chang* 122(1–2):185–199
- Bergot M, Cloppet E, Pérarnaud V, Déqué M, Desprez-Loustau ML (2004) Simulation of potential range expansion of oak disease caused by *Phytophthora cinnamomi* under climate change. *Glob Chang Biol* 10:1539–1552
- Bonet A, Pausas JG (2007) Old field dynamics on the dry side of the Mediterranean Basin: patterns and processes in semiarid SE Spain. In: Cramer VA, Hobbs RJ (eds) Old fields dynamics and restoration of abandoned farmland. Island Press, Washington, pp 247–264
- Bugalho MN, Caldeira MC, Pereira JS, Aronson J, Pausas JG (2011) Mediterranean cork oak savannas require human use to sustain biodiversity and ecosystem services. *Front Ecol Environ* 9:278–286
- Calkin DE, Thompson MP, Finney MA (2015) Negative consequences of positive feedbacks in US wildfire management. *For Ecosyst* 2:9
- Camia A, Liberta G, San-Miguel-Ayanz J (2017) Modeling the impacts of climate change on forest fire danger in Europe. JRC Technical Report <https://doi.org/10.2760/768481>
- Catry FX, Moreira F, Duarte I, Acacio V (2009) Factors affecting post-fire crown regeneration of cork oak (*Quercus suber*) trees. *Eur J Forest Res* 128:231–240
- Catry FX, Moreira F, Cardillo E, Pausas JG (2012a) Post-fire management of cork oak forests. In: Moreira F, Arianoutsou M, Corona P, De las Heras J (eds) Post-fire management and restoration of European forests. *Managing forest ecosystems*, vol 24. Springer, Berlin, pp 195–222
- Catry FX, Moreira F, Pausas JG, Fernandes PM, Rego F, Cardillo E, Curt T (2012b) Cork oak vulnerability to fire: the role of bark harvesting, tree characteristics and abiotic factors. *PLoS ONE* 7(6):e39810
- Chas-Amil ML, Touza J, García-Martínez E (2013) Forest fires in the wildland–urban interface: a spatial analysis of forest fragmentation and human impacts. *Appl Geogr* 43:127–137
- Corona P, Quatrini V, Schirru M, Dettori S, Puletti N (2018) Towards the economic valuation of ecosystem production from cork oak forests in Sardinia (Italy). *iForest* 11(5):660–667
- Costa A, Madeira M (2011) Assessment of Mediterranean evergreen oak (*Quercus suber* L.) woodlands loss. Consequence of climate changes effects? A case study at south-western Portugal. In: Proceedings of the global conference on global warming 2011. Lisbon, Portugal, 11–14 July
- Costa A, Madeira M, Plieninger T (2014) Cork oak woodlands patchiness: a signature of imminent deforestation? *Appl Geogr* 54:18
- Curt T, Frejaville T (2018) Wildfire policy in Mediterranean France: how far is it efficient and sustainable? *Risk Anal* 38(3):472–488
- Curt T, Pausas JG (2010) Are changes in fire regime threatening cork oak-shrubland mosaics? In: IUFRO-IALE international conference “landscapes forest and global change: new frontiers in management, conservation and restoration”. Bragança, Portugal, September 2010, pp 60–69
- Curt T, Borgniet L, Bouillon C (2013) Wildfire frequency varies with the size and shape of fuel types in south-eastern France: implications for environmental management. *J Environ Manag* 117:150–161
- Dettori S, Filigheddu MR (2003) La Sughericoltura. In: Minotta GF (ed) “L'Arboricoltura da Legno: un'attività produttiva al servizio dell'ambiente”. Avenue media, Bologna, pp 151–159
- Dettori S, Filigheddu MR (2016) Il sughero in enologia: analisi della filiera nazionale. *L'Ital Fore e Mont* 71(6):331–343
- Dettori S, Filigheddu MR, Deplano G, Escamilla Molgora J, Ruiu M, Sedda L (2018) Employing a spatio-temporal contingency table for the analysis of cork oak cover change in the Sa Serra region of Sardinia. *Sci Rep* 8:16946. <https://doi.org/10.1038/s41598-018-35319-1>
- Dimitrakopoulos AP, Vlahou M, Anagnostopoulou CG, Mitsopoulos ID (2011) Impact of drought on wildland fires in Greece: implications of climatic change? *Clim Chang* 109:331–347
- EEA (2007) European forest types. Categories and types for sustainable forest management reporting and policy. European Environment Agency (EEA) Technical Report no 9/2006, 2nd edn, Copenhagen
- EEA (2017a) Copernicus land monitoring service—Corine land cover. <https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-corine>
- EEA (2017b) Copernicus land monitoring service—EU-DEM. <https://www.eea.europa.eu/data-and-maps/data/copernicus-land-monitoring-service-eu-dem>
- DG Environment (2007) Interpretation manual of European Union habitats—EUR 27. p 142. (http://ec.europa.eu/environment/nature/legislation/habitatsdirective/docs/2007_07_im.pdf)
- FAO (2013) State of Mediterranean forests. p 174. <http://www.fao.org/docrep/017/i3226e/i3226e.pdf>
- Fernandes P, Rego FC (1996) Changes in fuel structure and fire behaviour with heathlands aging in Northern Portugal. In: 13th fire and forest meteorology conference. Lorne, Australia, pp 433–436
- Fernandes P, Davies GM, Ascoli D, Fernández C, Moreira F, Rigolot E, Stoof CR, Vega JA, Molina DM (2013) Prescribed burning in southern Europe: developing fire management in a dynamic landscape. *Front Ecol Environ* 11:e4–e14
- Ferreira-Leite F, Bento-Gonçalves A, Vieira A, Nunes A, Lourenço L (2016) Incidence and recurrence of large forest fires in mainland Portugal. *Nat Hazards* 84(2):1035–1053
- Fick SE, Hijmans RJ (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *Int J Climatol* 37:4302–4315
- Figueiredo E, Raschi A (2011) Un' immensa campagna avvolta dal verde. Reinventing rural areas in Italy through tourism promotional images. *Eur Countrys* 3(1):1–20
- Flohn, Fantechi (1984) The climate of Europe: Past, present and future. Springer, Berlin, p 357. ISBN:978-90-277-1745-0
- Forleo MB, Giaccio V, Giannelli A, Mastronardi L, Palmieri N (2017) Socio-economic drivers, land cover changes and the dynamics

- of rural settlements: Mt. Matese Area (Italy). *Eur Countrys* 9(3):435–457
- Hosmer DW, Lemeshow S (2000) *Applied logistic regression*, 2nd edn. John Wiley & Sons, New York. <https://doi.org/10.1002/0471722146>
- http://ec.europa.eu/environment/nature/natura2000/index_en.htm
- <http://www.sardegnageoportale.it/>
- <https://inventaire-forestier.ign.fr/>
- Ibáñez B, Gómez-Aparicio L, Ávila JM, Pérez-Ramos IM, Marañón T (2017) Effects of *Quercus suber* decline on woody plant regeneration: potential implications for successional dynamics in Mediterranean forests. *Ecosystems* 20:630
- Jiménez-Ruano A, Rodrigues Mimbreno M, de la Riva Fernández J (2017) Exploring spatial–temporal dynamics of fire regime features in mainland Spain. *Nat Hazards Earth Syst Sci* 17:1697–1711
- Kerckhof A, Spalevic V, Van Eetvelde V, Nyssen J (2016) Factors of land abandonment in mountainous Mediterranean areas: the case of Montenegrin settlements. *SpringerPlus* 5:485
- Koutsias N, Xanthopoulos G, Founda D, Xystrakis F, Nioti F, Pleniou M, Mallinis G (2012) On the relationships between forest fires and weather conditions in Greece from long-term 20 national observations (1894–2010). *Int J Wildland Fire* 22(4):493–507
- Kovats RS, Valentini R, Bouwer LM, Georgopoulou E, Jacob D, Martin E, Rounsevell M, Soussana J-F (2014) Europe. In: Barros VR, Field CB, Dokken DJ, Mastrandrea MD, Mach KJ, Bilir TE, Chatterjee M, Ebi KL, Estrada YO, Genova RC, Girma B, Kissel ES, Levy AN, MacCracken S, Mastrandrea PR, White LL (eds) *Climate change 2014: impacts, adaptation, and vulnerability. Part B: regional aspects. Contribution of working group II to the fifth assessment report of the intergovernmental panel on climate change*. Cambridge University Press, Cambridge, pp 1267–1326
- Levin N, Tessler N, Smith A, McAlpine C (2016) The human and physical determinants of wildfires and burnt areas in Israel. *Environ Manag* 58:549–562
- Lozano O, Salis M, Ager AA, Arca B, Alcasena FJ, Monteiro A, Finney MA, Del Giudice L, Scocciarro E, Spano D (2017) Assessing climate change impacts on wildfire exposure in Mediterranean areas. *Risk Anal* 37(10):1898–1916
- Luciano P, Roversi PF (2001) Fillofagi delle querce in Italia. *Industria Grafica Poddighe s.r.l, Sassari*, p. 161
- Luciano P, Lentini A, Cao OV (2005) La lutte aux lépidoptères défoliateurs des subéraies dans la Province de Sassari. *Industria Grafica Poddighe s.r.l, Sassari*
- Lung T, Lavalle C, Hiederer R, Dosio A, Bouwer LM (2013) A multi-hazard regional level impact assessment for Europe combining indicators of climatic and non-climatic change. *Glob Environ Chang* 23(2):522–536
- Madrigal J, Fernández-Migueláñez I, Hernando C, Guijarro M, Vega-Nieva DJ, Tolosana E (2016) Does forest biomass harvesting for energy reduce fire hazard in the Mediterranean basin? A case study in the Caroig massif (Eastern Spain). *Eur J For Res* 136:13–26
- Molina DM, Castellnou M, García D, Salgueiro A (2010) Improving fire management success through fire behavior specialists. In: *Towards integrated fire management—outcomes of the European Project Fire Paradox—EFI Research Report nr. 23. “Fire Paradox”, Project no. FP6-018505, European Commission*, pp 105–119. ISBN 970-952-5453-48-5. http://www.efi.int/files/attachments/publications/efi_rr23.pdf
- Montiel Molina C, Galiana-Martín L (2016) Fire scenarios in Spain: a territorial approach to proactive fire management in the context of global change. *Forests* 7:273
- Moreira F, Duarte I, Catry F, Acacio V (2007) Cork extraction as a key factor determining post-fire cork oak survival in a mountain region of southern Portugal. *For Ecol Manag* 253:30–37
- Moreira F, Viedma O, Arianoutsou M, Curt T, Koutsias N, Rigolot E, Barbati A, Corona P, Vaz P, Xanthopoulos G, Mouillot F, Bilgili E (2011) Landscape–wildfire interactions in southern Europe: implications for landscape management. *J Environ Manag* 92(10):2389–2402
- Moreno JM, Vazquez A, Velez R (1998) Recent history of forest fires in Spain. In: Moreno JM (ed) *Large forest fires*. Backhuys Publishers, Leiden, pp 159–185
- Moriondo M, Good P, Durao R, Bindi M, Giannakopoulos C, Cortes-Real J (2006) Potential impact of climate change on fire risk in the Mediterranean area. *Clim Res* 31:85–95
- Moritz MA, Batllori E, Bradstock RA, Gill AM, Handmer J, Hessburg PF, Leonard J, McCaffrey S, Odion DC, Schoennagel T, Syphard AD (2014) Learning to coexist with wildfire. *Nature* 515:58–66
- Oliveira S, Pereira JMC, San-Miguel-Ayanz J, Lourenço L (2014) Exploring the spatial patterns of fire density in Southern Europe using geographically weighted regression. *Appl Geogr* 51:143–157
- Oliveira V, Lauw A, Pereira H (2016) Sensitivity of cork growth to drought events: insights from a 24-year chronology. *Clim Chang* 137:261–274
- Oubrahim H, Boulmane M, Bakker, Augusto L, Halim M (2015) Carbon storage in degraded cork oak (*Quercus suber*) forests on flat lowlands in Morocco. *iForest* 9:125–137
- Pausas JG (1997) Resprouting of *Quercus suber* in NE Spain after fire. *J Veg Sci* 8:703–706
- Pausas JG (2004) Changes in fire and climate in the eastern Iberian peninsula (Mediterranean Basin). *Clim Chang* 63:337–350
- Pausas JG, Fernandez-Munoz S (2012) Fire regime changes in the Western Mediterranean Basin: from fuel-limited to drought-driven fire regime. *Clim Chang* 110:215–226
- Pausas JG, Paula S (2012) Fuel shapes the fire–climate relationship: evidence from Mediterranean ecosystems. *Glob Ecol Biogeogr* 21:1074–1082
- Pausas J, Vallejo VR (1999) The role of fire in European Mediterranean ecosystems. In: Chuvieco E (ed) *Remote sensing of large wildfires in the European Mediterranean basin*. Springer, Berlin, pp 3–16
- Pellizzaro G, Duce P, Ventura A, Zara P (2007) Seasonal variations of live moisture content and ignitability in shrubs of the Mediterranean Basin. *Int J Wildland Fire* 16:633–641
- Pereira H (2007) *Cork: biology, production and uses*. Elsevier Publishing, Amsterdam, p 336
- Pereira MG, Trigo RM, da Camara CC, Pereira JMC, Leite SM (2005) Synoptic patterns associated with large summer forest fires in Portugal. *Agric For Meteorol* 129:11–25
- Pereira MG, Calado TJ, DaCamara CC, Calheiros T (2013) Effects of regional climate change on rural fires in Portugal. *Clim Res* 57:187–200
- Piñol J, Beven K, Viegas D (2005) Modelling the effect of fire-exclusion and prescribed fire on wildfire size in Mediterranean ecosystems. *Ecol Model* 183:397–409
- Piñol J, Castellnou M, Beven KJ (2007) Conditioning uncertainty in ecological models: assessing the impact of fire management strategies. *Ecol Model* 207(1):34–44
- Plieninger TD, van der Horst D, Schleyer C, Bieling C (2014) Sustaining ecosystem services in cultural landscapes. *Ecol Soc* 19(2):59
- Rodrigues M, Jiménez-Ruano A, Peña-Angulo D, de la Riva J (2018) A comprehensive spatial-temporal analysis of driving factors of human-caused wildfires in Spain using geographically weighted logistic regression. *J Environ Manag* 225:177–192
- Romero-Calcerrada R, Perry GLW (2004) The role of land abandonment in landscape dynamics in the SPA ‘Encinares del río Alberche y Cofio, Central Spain, 1984–1999. *Landsc Urban Plann* 66:217–232

- Romero-Calcerrada R, Novillo J, Millington DA, Gomez-Jimenez I (2008) GIS analysis of spatial patterns of human-caused wildfire ignition risk in the SW of Madrid (Central Spain). *Landsc Ecol* 23(3):341–354
- Salis M, Ager AA, Arca B, Finney MA, Bacciu V, Duce P, Spano D (2013) Assessing exposure of human and ecological values to wildfire in Sardinia, Italy. *Int J Wildland Fire* 22(4):549–565
- Salis M, Ager AA, Finney MA, Arca B, Spano D (2014) Analyzing spatiotemporal changes in wildfire regime and exposure across a Mediterranean fire-prone area. *Nat Hazards* 71(3):1389–1418
- Salis M, Arca B, Alcasena F, Arianoutsou M, Bacciu V, Duce P, Duguay B, Koutsias N, Mallinis G, Mitsopoulos I, Moreno JM, Pérez JR, Rodríguez I, Xystrakis F, Zavala G, Spano D (2016a) Predicting wildfire spread and behavior in Mediterranean landscapes. *Int J Wildland Fire* 25(10):1015–1032
- Salis M, Laconi M, Ager AA, Alcasena FJ, Arca B, Lozano OM, Oliveira AS, Spano D (2016b) Evaluating alternative fuel treatment strategies to reduce wildfire losses in a Mediterranean area. *For Ecol Manag* 368:207–221
- Salis M, Del Giudice L, Arca B, Ager AA, Alcasena-Urdiroz F, Lozano O, Bacciu V, Spano D, Duce P (2018) Modeling the effects of different fuel treatment mosaics on wildfire spread and behavior in a Mediterranean agro-pastoral area. *J Environ Manag* 212:490–505
- San-Miguel-Ayanz J, Moreno JM, Camia A (2013) Analysis of large fires in European Mediterranean landscapes: lessons learned and perspectives. *For Ecol Manag* 294:11–22
- San-Miguel-Ayanz J, Durrant T, Boca R, Libertà G, Boccacci F, Di Leo M, López Pérez J, Schulte E (2016) Forest fires in Europe, Middle East and North Africa 2015. EUR 28158 EN. <https://doi.org/10.2788/914>. http://effis.jrc.ec.europa.eu/media/cms_page_media/40/Forest_fires_in_Europe_Middle_east_and_North_Africa_2015_final_pdf_JkX4Y13.pdf
- Santana J, Porto M, Reino L, Beja P (2011) Long-term understory recovery after mechanical fuel reduction in Mediterranean cork oak forests. *For Ecol Manag* 261:447–459
- Schaffhauser A, Curt T, Véla E, Taton T (2012) Fire recurrence effects on the abundance of plants grouped by traits in *Quercus suber* L. woodlands and maquis. *For Ecol Manag* 282:157–166
- Scozzafava S, De Sanctis A (2006) Exploring the effects of land abandonment on habitat structures and on habitat suitability for three passerine species in a highland area of central Italy. *Landsc Urban Plann* 75:23–33
- Silva JS, Catry F (2006) Forest fires in cork oak (*Quercus suber* L.) stands in Portugal. *Int J Environ Stud* 63:235–257
- Stephens SL, Collins BM, Biber E, Fulé PZ (2016) US federal fire and forest policy: emphasizing resilience in dry forests. *Ecosphere* 7:e01584
severicat.ctfc.cat
- Thompson MP, y Silva FR, Calkin DE, Hand MS (2017) A review of challenges to determining and demonstrating efficiency of large fire management. *Int J Wildland Fire* 26:562–573
- Thompson MP, MacGregor DG, Dunn CJ, Calkin DE, Phipps J (2018) Rethinking the wildland fire management system. *J For* 116:382–390
- Turco M, Bedia J, Di Liberto F, Fiorucci P, von Hardenberg J, Koutsias N, Llasat MC, Xystrakis F, Provenzale A (2016) Decreasing fires in Mediterranean Europe. *PLoS ONE* 11(3):e0150663. <https://doi.org/10.1371/journal.pone.0150663>
- Turco M, von Hardenberg J, Agha Kouchak A, Llasat MC, Provenzale A, Trigo RM (2017) On the key role of droughts in the dynamics of summer fires in Mediterranean Europe. *Sci Rep* 7:81
- Urbietta IR, Zavala MA, Marañón T (2008) Human and non-human determinants of forest composition in southern Spain: evidence of shifts towards cork oak dominance as a result of management over the past century. *J Biogeogr* 35:1688–1700
- Varela MC (2017) From genetic improvement to global change: what advantages and threats for the production, adaptation and evolution of cork oak? In: Dettori S, Fligheddu MR, Cillara M (ed) Proceedings of the international congress on cork oak trees and Woodlands, 25–26 May. Sassari (Italy). ISBN 978-88-907678-0-7
- Vega-Garcia C, Woodard T, Adamowicz WL, Lee B (1995) A logit model for predicting the daily occurrence of human caused forest fires. *Int J Wildland Fire* 5:101–111
- Vericat P, Piqué M, Koua O, Pla M (2010) Mapa de formacions forestals pures i mixtes de Catalunya a partir del Mapa Forestal de España 1:50.000 digitalizado. Centre Tecnològic Forestal de Catalunya, Solsona
- Viedma O, Moreno JM, Rieiro I (2006) Interactions between land use/land cover change, forest fires and landscape structure in Sierra de Gredos (Central Spain). *Environ Conserv* 33:212–222
- Viegas DX, Viegas MT (1994) A relationship between rainfall and burned area for Portugal. *Int J Wildland Fire* 4:11–16
- Viegas DX, Figueiredo Almeida M, Ribeiro LM, Raposo J, Viegas MT, Oliveira R, Alves D, Pinto C, Jorge H, Rodrigues A, Lucas D, Lopes S, Silva LF (2017) O complexo de incendios de Pedrogao Grande e concelhos limítrofes, iniciado a 17 de junho de 2017. <https://www.portugal.gov.pt/pt/gc21/comunicacao/documento/i40-complexo-de-incendios-de-pedrogao-grande-e-concelhos-limitrofes-iniciado-a-17-de-junho-de-2017>
- Vogiatzakis IN, Careddu MB (2003) Mapping the distribution and extent of *Quercus suber* habitats in Sardinia: a literature review and a proposed methodology. *Geogr Paper N. 171*, University of Reading, Reading UK. p 30
- Xystrakis F, Kallimanis AS, Dimopoulos P, Halley JM, Koutsias N (2014) Precipitation dominates fire occurrence in Greece (1900–2010): its dual role in fuel build-up and dryness. *Nat Hazards Earth Syst Sci* 14:21–32

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Affiliations

Michele Salis¹  · Bachisio Arca¹ · Fermin Alcasena-Urdiroz² · Antonella Massaiu³ · Valentina Bacciu⁴ · Frédéric Bosseur⁵ · Philippe Caramelle³ · Sandro Dettori⁶ · Ana Sofia Fernandes de Oliveira⁶ · Domingo Molina-Terren⁷ · Grazia Pellizzaro¹ · Paul-Antoine Santoni⁵ · Donatella Spano^{4,6} · Cristina Vega-Garcia^{2,8} · Pierpaolo Duce¹

Bachisio Arca
b.arca@ibimet.cnr.it

Fermin Alcasena-Urdiroz
ferminalcasena@eagrof.udl.cat

Antonella Massaiu
antonella.massaiu@onf.fr

Valentina Bacciu
valentina.bacciu@cmcc.it

Frédéric Bosseur
bosseur@univ-corse.fr

Philippe Caramelle
philippe.caramelle@onf.fr

Sandro Dettori
sdettori@uniss.it

Ana Sofia Fernandes de Oliveira
acortez@uniss.it

Domingo Molina-Terren
dmolina@pvcf.udl.es

Grazia Pellizzaro
g.pellizzaro@ibimet.cnr.it

Paul-Antoine Santoni
santoni@univ-corse.fr

Donatella Spano
donatella.spano@cmcc.it; spano@uniss.it

Cristina Vega-Garcia
cvega@eagrof.udl.cat

Pierpaolo Duce
p.duce@ibimet.cnr.it

- ¹ National Research Council (CNR), Institute of Biometeorology (IBIMET), Traversa La Crucca 3, 07100 Sassari, Italy
- ² Department of Agricultural and Forest Engineering, University of Lleida, Avenida da Rovira Roure 191, 25198 Lleida, Spain
- ³ Office National des Forêts (ONF), Résidence la Pietrina, Avenue de la Grande Armée, 20000 Ajaccio, France
- ⁴ IAFES Division, Euro-Mediterranean Center on Climate Change (CMCC), Via De Nicola 9, 07100 Sassari, Italy
- ⁵ SPE-UMR 6134, CNRS, University of Corsica, Campus Grossetti, B.P. 52, 20250 Corte, France
- ⁶ Department of Agriculture, University of Sassari, Via De Nicola 9, 07100 Sassari, Italy
- ⁷ Department of Crop and Forest Science, University of Lleida, Avenida da Rovira Roure 191, 25198 Lleida, Spain
- ⁸ Forest Sciences Centre of Catalonia (CTFC), Carretera de St. Llorenç de Morunys, km 2, 25280 Solsona, Spain